



PROCEEDINGS  
OF THE  
**ROYAL SOCIETY OF VICTORIA**



VOLUME 115 NUMBER 1/2

ROYAL SOCIETY'S HALL

9 VICTORIA STREET

VICTORIA 3000 AUSTRALIA

30 NOVEMBER 2003

MELBOURNE

# ESTABLISHMENT OF THE ROYAL SOCIETY OF VICTORIA

The Publications of the Royal Society of Victoria and of the  
Societies amalgamated with it

---

VICTORIAN INSTITUTE FOR THE ADVANCEMENT OF SCIENCE  
Transactions and Proceedings, Vol. 1, 1854/55.

PHILOSOPHICAL SOCIETY OF VICTORIA  
Transactions, Vol. 1, 1854/55.

*These two Societies then amalgamated and became—*

PHILOSOPHICAL INSTITUTE OF VICTORIA  
Transactions, Vol. 1-4, 1854/56-1859.

*The Society then became—*

ROYAL SOCIETY OF VICTORIA  
Transactions and Proceedings (Vol. 5, entitled Transactions), Vol. 5-24, 1860=1887.  
Transactions, Vol. 1, 2, 3 (Pt 1 only was published), 4, 5 (Pt 1 only), 6, 1888-1914.  
Proceedings (New Series), Vol. 1-82, 1888-1969.  
Proceedings, Vol. 83 ... 1969 ...  
(*'New Series' deleted from Title Page*)

MICROSCOPICAL SOCIETY OF VICTORIA  
Journal (Vol. 1, Pt 1, entitled Quarterly Journal), Vol. 1 (Pt 1 to 4), 2 (Pt 1), title page and index [all  
published], 1879-82.  
[*The society then combined with the ROYAL SOCIETY OF VICTORIA*]  
*Note—Most of the volumes published before 1890 are out of print.*

---

Vol. 46, Part 2, contains an author index of the Royal Society of Victoria and of the Societies amalgamated with it from 1855-1934.

Vol. 73 contains an author index of the Proceedings of the Royal Society of Victoria, Vol. 47 (1934/5)-70 (1958).

Vol. 84, part 2, contains an author index of the Proceedings of the Royal Society of Victoria, Vol. 71 (1959)-84 (1971).

Vol. 99, Part 4, contains an author index of the Proceedings of the Royal Society of Victoria, Vol. 85 (1972)-98 (1986).

Vol. 113, Number 1, contains an author index of the Proceedings of the Royal Society of Victoria, Vol. 99 (1987)-112 (2000).

*Many of the above volumes are available for purchase. All orders should be sent to:*


*The Executive Officer,  
Royal Society of Victoria,  
9 Victoria Street,  
Melbourne, Victoria 3000, Australia  
Tel: (03) 9663 5259  
Fax: (03) 9663 2301  
Email address: rsvine@vnet.net.au*



PROCEEDINGS  
OF THE  
ROYAL SOCIETY OF VICTORIA  
INCLUDING  
TRANSACTIONS OF MEETINGS

Volume 115

NUMBERS 1/2



ROYAL SOCIETY'S HALL  
9 VICTORIA STREET, MELBOURNE, VICTORIA 3000

This issue is sponsored by Deakin University and the Deakin University Global Change Research group



and the Department of Natural Resources and Environment

Natural Resources and Environment  The Place To Be

## CONTENTS OF VOLUME 115, NUMBER 1/2

### PROCEEDINGS

- B. MACKNESS 1  
A review of nematophagy by terrestrial tardigrades with new observations on some Australian species.
- BASDEN, ALISON 7  
Early Devonian (Emsian) vertebrate microremains from the Buchan Group, Victoria, Australia
- FRAMENAU, VOLKER W. 27  
Two Alpine Wolf Spiders of Australia: *Artoria alta* sp. nov., and the male of *Lycosa musgravei* McKay, 1974 (Araneae, Lycosidae)
- GERDTZ, W. R. & ARCHIBOLD, N. W. 35  
*Glancodon ballaratensis* (Marsupialia, Dasyuridae), a late Pliocene 'devil' from Batesford, Victoria
- GERDTZ, W. R. & ARCHIBOLD, N. W. 45  
An early occurrence of *Sarcophilus laniarius harrisii* (Marsupialia, Dasyuridae) from Nelson Bay, Victoria
- YEN, ALAN L. & WAINER, JOHN 55  
Scolopendrid centipedes caught by pitfall trapping in the Uluru-Kata Tjuta National Park, Northern Territory, including a record of an apparent 'outbreak'
- BIRCH, W. D. & SAMUELS, L. E. 67  
The Ballarat meteorite, a fossil IAB iron from Victoria, Australia

### TRANSACTIONS

- LAW, PHILLIP i  
Lecture to the Danish Club, 18<sup>th</sup> September, 2003

## EDITORIAL POLICY

The RSV disseminates new scientific knowledge and encourages scientific debate through the medium of its international journal *Proceedings of the Royal Society of Victoria*.

Each volume of the 'Proceedings' may consist of two sections:

The 'Proceedings' proper consists of peer reviewed contributions. Each contribution is reviewed by two reviewers and, in the cases of widely divergent reviewers opinions, a third reviewer may be consulted. The editor's decision, through the Editorial Committee, as to the acceptance or rejection of a submitted or reviewed contribution, is final.

The 'Transactions' of the Society consists of unreviewed items of scientific interest such as occasional addresses, articles of debate or discussion or other contributions deemed appropriate by the editor and in the public's interest. The responsibility of the content of such contributions rests solely with the author.

The RSV invites contributions for the *Proceedings* from authors across the various disciplines of the physical and natural sciences and on issues concerning technology and the applied sciences.

Chief Editor: Montague Grover

PROCEEDINGS  
OF THE  
ROYAL SOCIETY OF VICTORIA  
PEER REVIEWED PAPERS





# A REVIEW OF NEMATOPHAGY BY TERRESTRIAL TARDIGRADES WITH NEW OBSERVATIONS ON SOME AUSTRALIAN SPECIES

BRIAN MACKNESS

School of Biological Sciences, University of New South Wales, New South Wales 2054, Australia

Current address: PO Box 560, Beerwah, Queensland 4519, Australia

MACKNESS, B.S., 2003:06:15. A review of nematophagy by terrestrial tardigrades with new observations on some Australian species. *Proceedings of the Royal Society of Victoria* 115(1): 1-5. ISSN 0035-9211.

A literature review of nematophagy by terrestrial tardigrades was undertaken including a survey of 93 species of the genus *Macrobiotus* to determine whether there were any morphological criteria common to all nematophagous species. Preliminary results suggest that all nematophagous species have buccal tube widths greater than 17% of the tube length and that they also possess peribuccal lamellae. Two new nematophagous tardigrades, *Macrobiotus hieronimi* Pilato & Claxton, 1988 and *M. peteri* Pilato et al., 1989 have been recognized as well as further records of nematode predation by tardigrades from Australia.

**Key words:** Tardigrades, nematophagy, buccal tubes, peribuccal lamellae

TERRESTRIAL tardigrades are primarily vegetarian, although some are known to take animal food including nematodes (Ramazzotti & Muccii 1983). There are few reports of nematophagy however found in the literature. Linford & Oliveira (1938) reported three predacious tardigrades (*Macrobiotus* spp.) attacking root-knot nematodes *Heterodera* sp. (now various species of *Meloidogyne*). Le Gros (1958) stated that tardigrades attacked nematodes in addition to rotifers and other tardigrades. Hutchinson & Streu (1960) reported an unidentified tardigrade feeding on the nematodes *Trichodorus* sp. and *Tylenchus* sp. in the United States and in Sri Lanka. Doneaster & Hooper (1961) and Doneaster (1962) recorded *Macrobiotus richtersi* Murray, 1911 piercing nematodes and suggested that a toxin might have been secreted to immobilize large nematodes such as mononchids. Esser (1963) and Esser & Sobers (1964) examined the possibility of tardigrades being used as biological control agents of nematodes. Macrobiotid tardigrades, in citrus orchards, have been recorded feeding on the nematodes *Tylenchulus* sp. and *Meloidogyne* sp. (Boosalis & Manakau 1965, Stirling & Mankau 1977).

Sayre (1969) cultured the tardigrade *Hypsibius* (= *Isolytsibius*) *myrops* (Du Bois-Reymond Mareus, 1944) on nematodes for several months. He also observed individuals feeding on the plant nematodes *Meloidogyne* sp. and *Ditylenchus* sp. Hallas & Yeates (1972) recorded *Macrobiotus harmsworthi* Murray, 1907 feeding on nematodes and reported a strong

correlation between total number of nematodes and monthly tardigrade populations. Hallas & Yates (1972) also suggested that *Hypsibius* (*I.*) *myrops* (*sensu* Sayre 1969) might be identical to *Hypsibius* (*I.*) *prosostomus*, which they also reported feeding on nematodes.

Several other authors have reported general incidents of predation by tardigrades (Cayrol 1976, Morgan 1977, Small 1988). Further investigations of predation by tardigrades were undertaken by Sayre & Wergin (1979) using a scanning electron microscope. Esser (1990) summarized more than 25 years of observing nematodes and tardigrades. These observations primarily concerned hypsibiid tardigrades feeding on a variety of nematodes. Bird (1996) figured *Macrobiotus* cf. *pseudohufelandi* Iharos, 1966 eating an unidentified soil nematode.

Hallas & Yeates (1972: 289) characterized predacious tardigrades in having '... a mouth opening (oral aperture) which points directly forward, a short, wide pharyngeal tube, strongly built stylets and a large bulbus'. A survey was undertaken of published buccal tube widths of 93 species of *Macrobiotus* to ascertain whether this criterion could be used as a guide to recognizing predacious tardigrades. The incidence of buccal lamellae was also examined. The results of this survey are reported in Table 1. The author reports two hitherto unknown species of nematophagous tardigrades as well as further records of nematode predation by tardigrades from Australia.

Species	BTR	Length	Lamellae
<i>M. beotiae</i> Durante & Maucci, 1979	(25% of length)	880 µm	Yes
<i>M. gildae</i> Maucci & Durante Pasa, 1980	(22% of length)	500 µm	Yes
<i>M. grandis</i> Richters, 1911	(17% of length)	900 µm	Yes
<i>M. harmsworthi</i> Murray, 1907	(20% of length)	650 µm	Yes
<i>M. hieronimi</i> Pilato & Claxton, 1988	(17% of length)	488 µm	Yes
<i>M. maucci</i> Pilato, 1974	(22% of length)	450 µm	Yes
<i>M. petersi</i> Pilato et al. 1989	(17% of length)	690 µm	Yes
<i>M. pseplus</i> du Bois-Reymond Marcus, 1944	(17% of length)	800 µm	Yes
<i>M. pseudohufelandi</i> Iharos, 1966	(18% of length)	510 µm	Yes
<i>M. recens</i> Cuenot, 1932	(21% of length)	624 µm	Yes
<i>M. richtersi</i> Murray, 1911	(20% of length)	750-1000 µm	Yes
<i>M. snaresensis</i> Horning et al., 1978	(22% of length)	450 µm	Yes
<i>M. spectabilis</i> Thulin, 1928	(17% of length)	700 µm	Yes

Table 1. Species of *Macrobiotus* with buccal tube widths greater than 17% of buccal tube length. (Source Ramazzotti & Maucci 1983 and references cited in text). Known nematophagous tardigrades highlighted in bold. (BTR = Buccal tube ratio)

### Feeding by tardigrades

Tardigrades have a complex series of organs associated with feeding. These include the buccal orifice, stylets, the pharynx and the oesophagus. The oral aperture can be terminal as in most macrobiotids or slightly ventral as in most hypsibids. The buccal cavity opens with a circular ring that may support a crown of buccal lamellae (Fig 1A). These allow the mouth to be applied like a sucker to prey (Ramazzotti & Maucci 1983). There are piercing organs on either side of the mouth called stylets. Their composition remains unclear although Marcus (1927) and Kaestner (1968) suggest they are calcareous. Bird (1996) has recorded marked bi-refringence of the stylets under polarized light and suggested this may be due to their inferred crystalline nature. The stylets themselves, are operated by several retracting and protracting muscles. Tardigrades use these stylets to pierce the cuticles of their prey. The buccal cavity is followed by a buccal tube that extends into the pharynx. The pharynx functions as a sucking organ and is generally ovoid in shape (Ramazzotti & Maucci 1983). Food is not masticated but passed by sucking action through a short oesophagus into the midgut where digestion takes place.

In some tardigrades, particularly Family Macrobiotidae, there is a series of circular swellings called placoids in the pharyngeal bulb (Fig 1B). They are thought to play a role in mechanical reinforcement of the pharynx (Ramazzotti & Maucci 1983). There

are also large rostral or salivary glands located lateral to the buccal tube and pharyngeal bulb. It has not been confirmed whether these glands secrete digestive juices and indeed some authors, such as Cuénot (1932), have ascribed an excretory function to them.

There appears to be no pattern to the choice or size of nematode taken as prey by tardigrades (Doncaster & Hooper 1961). Tardigrades have been recorded passing within 10 µm of a nematode without any attempt to attack (Esser 1990). Likewise, not all attacks by tardigrades are successful. Esser (1990) observed an attack by a tardigrade on the nematode *Enchodelus* (three and a half times larger than the tardigrade) that was unsuccessful as well as another on a large dorylaim. Prey is normally pierced with the stylets and the body contents sucked out. Several tardigrades may attack the same nematode simultaneously (Doncaster & Hooper 1961).

Sometimes the prey is swallowed whole. Ramazzotti & Maucci (1983) observed *Macrobiotus richtersi*, with a buccal tube of 12 µm, with a whole nematode inside its stomach. Hallas & Yeates (1972) observed *M. harmsworthi* with a nematode tail projecting out of its cloaca. Apparently it had ingested the nematode tail first and the thinner posterior end had passed through the gut unruptured while the anterior end had been digested.

When tardigrade attacks occur, they can be quite rapacious and sustained. Esser & Sobers (1964: 333) reported an earlier observation by Esser of tardigrades






Fig. 1. A, Peribuccal lamellae at entrance to buccal cavity. B, Pharyngeal bulb with placoids. C, *Macrobionus* sp. from Mt Ainsley, ACT with nematode. Scale bar = 100  $\mu$ m

'viciously attacking nematodes, ripping many of them to pieces and swallowing some of them whole'. Hutchinson & Streu (1960) reported that tardigrades made distinct tears in the cuticles of nematodes while Esser (1990) recorded a tardigrade holding on to a nematode for over three hours. Doncaster & Hooper

(1961) suggested that tardigrades might inject a toxic secretion into nematodes in order to subdue them.

The following tardigrade species from Australia and Lord Howe Island have been observed feeding on nematodes.

*Macrobotus hieronimi* Pilato & Claxton, 1988

One specimen (LH3-13) collected by Jo Mon from Lord Howe Island (6/7/80) and mounted by S.K. Claxton. Tardigrade is 420 µm long and recovered from damp moss *Ptychomitrium* sp. with 25-50% moisture content. The nematode is being attacked head on and all of the pharyngeal region has been consumed and internal structures dissolved. Consequently identification of the nematode is not possible. This is the first published record of nematophagy by this species.

*Macrobotus peteri* Pilato et al., 1989

Three specimens (ATS1065-3, 1067-16, 1090-10) collected and mounted by S.K. Claxton, Douglas Park, NSW. Tardigrades found in moss *Dicnemoloma* sp. The nematodes cannot be identified. Tardigrades are 320 µm, 420 µm and 430 µm long respectively. This is the first published record of nematophagy by this species.

*Macrobotus richtersi* Murray, 1911

One specimen (ATS 417-1) collected and mounted by S.K. Claxton, Cedarvale, NSW from moss/lichen. The nematode cannot be identified. The tardigrade is 420 µm long.

*Milnesium tardigradum* Doyère, 1840

One specimen (ATS 1058-37) collected by S.K. Claxton, Cambewarra Mountains, NSW. Tardigrade found in foliose lichen *Parmotrema* sp. and is 550 µm long. The tardigrade has attacked the nematode (which appears to be a female enopliid) in the middle of the body. It has a characteristic cylindrical pharynx, the posterior portion of which has been partly digested (A. Bird pers. comm.). *Milnesium tardigradum* is an active predator of both nematodes and other tardigrades.

*Macrobotus* sp. (Fig 1C)

One specimen (ATS 507/13) collected and mounted by G. Carruthers, east face of Mt Ainsley, ACT, 10 metres below summit. The tardigrade was found in wet moss and lichen. It is holding the nematode in its mouth.

The body walls of the nematode are constricted at either end of the tardigrade's mouth. The nematode can not be identified.

## DISCUSSION

Four genera of tardigrades are known to prey on nematodes - *Milnesium*, *Macrobotus*, *Isohypsibius* and *Hypsibius*. Six species of *Macrobotus* have been recorded feeding on nematodes. Preliminary results of a survey of 93 macrobiotid tardigrades suggests that all nematophagous species have proportionally wider buccal tubes with widths at least 17% of buccal tube length. They also all possess buccal lamellae. Using these criteria, a further eight tardigrades may be potentially nematophagous. These are *Macrobotus beotiae* Durante & Maueci, 1979; *M. gildae* Maueci & Durante Pasa, 1980; *M. grandis* Richters, 1911; *M. maucci* Pilato, 1974; *M. psephus* du Bois-Reymond Mareus, 1944; *M. recens* Cuénot, 1932; *M. suaresensis* Horning et al., 1978 and *M. spectabilis* Thulin, 1928. Further field collecting of these potential nematophagous species is needed to test the hypothesis of Hallas & Yeates (1972) that there is a morphological basis for nematode predation by tardigrades. The width of the buccal tube is not correlated to the length of the tardigrade (Ramazzotti & Maueci 1983) nor does there seem to be any relationship between the size of the nematode being attacked and the size of the attacking tardigrade with nematodes three times larger being attacked (Esser 1990). Large vigorous nematodes may be able to squirm free although observations show that tardigrades can be quite dogged in their attempts to subdue their prey often hanging on for hours (Esser 1990).

It appears that there are many factors involved in nematode prey selection by tardigrades. Hallas & Yeates (1972) have shown a direct correlation between numbers of *Macrobotus harmsworthi* and the availability of prey, even when environmental factors are taken into account (simple rank correlation +0.273; moisture excluded = 0.287; temperature excluded +0.286). When Esser (1990) placed several large populations of tardigrades on water agar containing numerous nematodes of many genera, no feeding or attacking was noted for several weeks. Esser (1990) suggested that the rarity of attacks in agar solution was because the conditions under which tardigrades feed are rarely obtained outside their normal habitat. There also appears to be no correlation between what types of nematodes are eaten by tardigrades. Bacteriophagous, phytophagous and a variety of predacious nematodes have all been taken. One of the problems



in ascertaining what types of nematodes become prey is that the feeding action of tardigrades normally dissolves vital taxonomic features of the nematodes concerned. Tardigrade/nematode interactions occur in a variety of microhabitats and under a range of environmental conditions. Further research needs to be undertaken to understand the roles that these various factors play in both prey selection by tardigrades and the frequency and cause of attacks.

#### ACKNOWLEDGEMENTS

I wish to thank Glen Carruthers, Melbourne and Sandra Claxton, Macquarie University who made specimens available for this study. Donald Horning, Tumblegum Research Laboratory, Loomberah, New South Wales, Australia and the late Alan Bird, Mitcham, South Australia provided encouragement to publish the work as well as commenting on various drafts of the manuscript.

#### REFERENCES

- BIRD, A.F., 1996. Studies on the soil-inhabiting tardigrade, *Macrobiotus* cf. *psendohmfelandi*, from South Australia. *Transactions of the Royal Society of South Australia* 120(4): 147–154.
- BOOSALIS, M.G. & MANKAU, R., 1965. Parasitism and predation of soil microorganisms. In *Ecology of Soilborne Plant Pathogens*, K.F. Baker & W.C. Snyder, eds., University of California Press, Berkeley, 374.
- CAYROL, J.C., 1976. Relations ecologiques entre Nematodes et autres organismes terrioles. *Bulletin de la Societe Zoologique de France* 101: 872.
- CUÉNOT, L., 1932. Tardigrades. In *Faune de France*, Volume 24, Paul Lechevalier, ed. Paris, 1–96.
- DONCASTER, C.C., 1962. Predators of soil nematodes (film) *Parasitology* 52, 19 pp.
- DONCASTER, C.C. & HOOPER, D.J., 1961. Nematodes attacked by protozoa and tardigrades. *Nematologica* 6: 333–335.
- ESSER, R.P., 1963. Nematode interactions in plates of non-sterile water agar. *Soil Crop Science Society of Florida* 23: 121–138.
- ESSER, R.P., 1990. Tardigrades attacking nematodes. Nematology Circular No. 177 (Contribution No. 396, Bureau of Nematology). Florida Department of Agriculture and Consumer Services. Division of Plant Industry. 4 pp.
- ESSER, R.P. & SOBERS, E.K., 1964. Natural enemies of nematodes. *Soil Crop Science Society of Florida* 24: 326–353.
- HALLAS, T.E. & YEATES, G.W., 1972. Tardigrada of the soil and litter of a Danish beech forest. *Pedobiologia* 12: 287–304.
- HUTCHINSON, M.T. & STREU, H.T., 1960. Tardigrades attacking nematodes. *Nematologica* 5: 149.
- KAESTNER, A., 1978. *Invertebrate Zoology*. Vol II. John Wiley & Sons Inc, New York.
- LE GROS, A.E., 1958. How to begin to study tardigrades. *Countryside* 18: 322.
- LINFORD, M.B. & OLIVEIRA, J.M., 1938. Potential agents of biological control of plant parasitic nematodes. *Phytopathology* 28: 14.
- MARCUS, E., 1927. Zur Anatomie und Ökologie mariner Tardigraden. *Zoologische Jahrbücher Abteilung für Systematik Ökologie und Geographie der Tiere* 53: 487–558.
- MORGAN, C.I., 1977. Population dynamics of two species of Tardigrada, *Macrobiotus hmfelandi* (Schultz) and *Echiniscus (Echiniscus) testudo* (Doyère) in roof moss from Swansea. *Journal of Animal Ecology* 46: 263–279.
- RAMAZZOTTI, G. & MAUCCI, E., 1983. Il Phylum Tardigrada. (III edizione riveduta e aggiornata). *Memoire dell'Istituto Italiano di Idrobiologia Dott. Marco De Marchi, Pollanza* 41: 1–1012.
- SAYRE, R.M., 1969. A method of culturing a predaceous tardigrades on the nematode *Panagrellus redivivus*. *Transactions of the American Microscopical Society* 88(2): 266–274.
- SAYRE, R.M. & WERGIN, W.P., 1979. The use of SEM to classify and evaluate the parasites and predators of pest nematodes. *Scanning Electron Microscopy* 1979(3): 89–96.
- SMALL, R.W., 1988. Invertebrate predators. pp. 73–92 in *Diseases of nematodes* Vol 2. ed. by G.O. Poinar Jr and H. -B. Jansson. CRC Press Inc., Boca Raton Florida.
- STIRLING, G.R. & MANKAU, R., 1977. Biological control of nematode parasites of citrus by natural enemies. *Proceedings of the International Society of Citriculture* 3: 843.

Manuscript received 1 August 1999

Revision accepted 8 January 2003



# EARLY DEVONIAN (EMSIAN) VERTEBRATE MICROREMAINS FROM THE BUCHAN GROUP, VICTORIA, AUSTRALIA

ALISON BASDEN

Centre for Ecostratigraphy and Palaeobiology, Department of Earth and Planetary Sciences, Macquarie University, New South Wales 2109, Australia

BASDEN, A., 2003:06:15. Early Devonian (Emsian) vertebrate microremains from the Buchan Group, Victoria, Australia. *Proceedings of the Royal Society of Victoria* 115(1) : 7-26. ISSN 0035-9211.

The vertebrate microfauna from the Early Devonian Buchan Group in northeastern Victoria is dominated by remains of acanthodians and placoderms, with osteichthyans less common, and chondrichthyans and thelodonts rare. Acanthodian remains include scales of *Nostolepoides platymarginata*, "*Nostolepis*" *guangxiensis*, *Gomphonchus*? *bogongensis*, scales similar to those of *Cheiracanthoides comptus* and *C. wangi*, and possible *Rhadinacanthus* scales. Scales of arthrodire *Goodradigbeeon* and *Buchanosteus* are the most abundant placoderm remains; placoderm plate fragments are attributed to *Buchanosteus* and *Murrindalaspis*. Osteichthyan remains include a scale of the actinopterygian *Ligulalepis toombsi*, and onychodont scales and teeth. Taxa previously undescribed from Buchan are acanthodians "*Nostolepis*" *guangxiensis*, *Cheiracanthoides* sp. cf. *C. wangi* and *Rhadinacanthus*, chondrichthyan *Ohirolepis* sp., and osteichthyan *Ligulalepis toombsi*.

**Key words:** Early Devonian, Emsian, vertebrate microfossil, acanthodian, placoderm, chondrichthyan, osteichthyan

DEVONIAN fossil fish remains have been known from the Buchan area of northeastern Victoria for well over a century. McCoy (1876) described as '*Asterolepis ornata* var. *australis*' a placoderm bone that has subsequently been shown to be portion of the skull of a large arthrodire (Young 1979). From the same area, an almost complete arthrodire skull was initially described by Chapman (1916), subsequently prepared in hydrochloric acid by Hills (1936) who referred it to the European genus *Coccosteus*, and eventually assigned to the new genus *Buchanosteus* by Stensiö (1945). Hills (1936) also described a mandible of the lungfish *Dipnorhynchus* from the same area.

The only recent description of macrovertebrate material from Buchan is by Long (1984a), who assigned various disarticulated plates from the McLarty Member of the Murrindal Limestone to *Buchanosteus confertituberculatus*, and described two new acanthothoracids, *Murrindalaspis wallacei* and *M. bairdi*. He identified several forms first described from the coeval Taemas/Wee Jasper fauna in southern New South Wales: brachythoracids *Arenipiscis westolli*, *Errolosteus* sp. cf. *E. goodradigbeensis*, and *Taemasosteus mclartiensis*, and petalichthyid *Wijdeaspis warrooensis*. Long (1984b) illustrated another *Buchanosteus* specimen, which included scales described by Burrow & Turner (1998, 1999). The only

other vertebrate microremains known from the Buchan region are scales of the dipnoan *Dipnorhynchus* illustrated by Thomson & Campbell (1971, figs 86-88).

The Early Devonian Buchan Group at Buchan, Victoria (Emsian: *dehiscens* - *serotinus* zones) comprises three formations (Fig. 1). The Buchan Caves Limestone, predominantly calcarenites and calcisiltites with dolomites and dolomitic limestones (*dehiscens* Zone) is overlain by the nodular and impure limestones and shales of the Taravale Formation (*dehiscens* - *serotinus* zones) and the well-bedded and massive limestones with subordinate mudstones of the Murrindal Limestone (*perboux* Zone). Detailed geology and conodont biostratigraphy were discussed by Mawson (1987) and Mawson et al. (1988, 1992).

Other studies have demonstrated the generally fossiliferous nature of these horizons, describing an invertebrate fauna of abundant stromatoporoids and corals associated with brachiopods, bivalves and ostracods in the Buchan Caves Limestone (Webby et al. 1993; Talent 1995), and more diverse faunas of brachiopods, corals, and trilobites, the oldest known ammonoids, daeryoeonarids, ostracods, agglutinated foraminiferans, seolecodonts and chitinozoans in the overlying Taravale Formation (Teichert & Talent 1958; Mawson 1987; Winchester-Seeto 1996).






Fig. 1. Geology of Buchan Group around Buchan and The Basin, eastern Victoria (modified from Mawson et al. 1992 and Winchester-Secto 1996).



Bed-by-bed sampling of long stratigraphic sections through the Buchan Group (Mawson 1987; Mawson et al. 1988, 1992) has resulted in acetic acid-insoluble residues containing both conodonts and microvertebrates. Identifiable microvertebrate remains occurred in three sections (Slocombe's - SL/SLO, 42 samples; McLarty's - Mc, 120 samples; and Gelantipy Road - G.Rd, 80 samples), spanning the *dehiscens*, *perbonus* and *inversus* zones. The turiniid thelodont scales from these sections, referred to *Turinia* sp. cf. *T. australiensis*, have been published elsewhere (Basden 1999a). This paper describes and figures the remaining microvertebrate taxa. The present study has increased the known geographic ranges of acanthodians "*Nostolepis*" *guangxiensis*, *Cheiracanthoides* sp. cf. *C. wangi* and *Rhadinacanthus*, chondrichthyan *Ohiolepis* sp., and osteichthyan *Ligulalepis toombsi*, all of which are reported for the first time from Buchan.

In the systematic descriptions below, the following abbreviations are used for measured stratigraphic sections: **SL/SLO**: Slocombe's section at The Basin, near Buchan, Buchan Caves Limestone and Taravale Formation, *?pirenae-dehiscens-perbonus* zones; **Mc**: McLarty's section near Buchan, Pyramids Member of Taravale Formation and Murrindal Limestone, *?dehiscens-perbonus* zones; **G.Rd**: Gelantipy Road section at Buchan, Taravale Formation, *perbonus-inversus* zones (See Fig. 1).

Samples are identified by the section prefix followed by the true height in metres above base of section (indicated by "m"). This is to distinguish between the true height above base of section, used in this study, and the sample number given by Mawson (1987), Mawson & Talent (1994), and Mawson et al. (1988, 1992) at the time of collection, which indicated measurement along the ground. Figured specimens are held in the palaeontology collections of the Australian Museum, Sydney, and bear the prefix AMF.

## SYSTEMATIC PALAEOLOGY

Superclass **Gnathostomata** Gegenbauer 1874

Class **Placodermi** M'Coy 1848

Order **Acanthothoraci** Stensiö 1944

Family **Weejasperaspididae** White 1978

Genus **Murrindalaspis** Long 1984

**Diagnosis** (Long 1984a; Long & Young 1988). Dermal ornament of short crenulate ridges, approximately one-third as long as broad, and zones of stout polygonal or stellate tubercles. Body scales

rhombic with anterior overlap flanges; exposed area with similar ornament to dermal bones.

### **Murrindalaspis** sp.

Fig. 2A

**Material.** AMF101198 from Mc41.5-42.7m (Pyramids Member of Taravale Formation, *dehiscens* or *perbonus* Zone), plus seven other scales and fragments from Mc3.4-3.6m, 8.5m, 18.6-21.3m, 27.7-30.5m, 41.5-42.7m (Pyramids Member of Taravale Formation, *dehiscens* or *perbonus* Zone), and Mc156.1m, 180.1m (Murrindal Limestone, *perbonus* Zone).

**Remarks.** Long (1984a) erected the genus from medial dorsal plates recovered from the Murrindal Limestone (*perbonus* Zone) near Buchan, Victoria. The only other described specimens from southeastern Australia are a sclerotic capsule and associated plates and body scales from the *Receptaculites* Limestone near Taemas (Long & Young 1988:74, figs 9, 10) and three body scales from the same sample (Burrow & Turner 1998:687, figs 10A-E), and a preliminary report of similar scales or plate fragments from the Lochkovian-Pragian Garra Limestone in central NSW, the Murrindal Limestone, and the upper Cavan Formation (*dehiscens* Zone) at Taemas (Basden et al. 2000, fig. 7.1).

**Description and Discussion.** The scale crowns and plate fragments are ornamented with elongate or stellate tubercles with radiating ridges that appear nodular under high magnification. Long & Young (1988) described the elongate tubercles on the plates as being 3 to 7 times as long as wide, and Burrow & Turner (1998) specified 4-20 nodose radiating ridges on each tubercle.

The scales are generally rhombic or subcircular, with diameter 1-2mm, and a thin margin that can be either faintly radially grooved or granular-textured. Neighbouring scales overlap this thin basal margin (Long & Young 1988), which can extend all around the scale, or be confined to the anterior and lateral margins. The base varies from deeply concave with openings into the tubercles, to gently concave with no canal openings, to flat. Tubercle length varies, with central elongate tubercles, aligned antero-posteriorly, surrounded by 3-12 smaller or more circular tubercles. Commonly, anteriormost tubercles tend to be lower and rounded, and those at the posterior flatter and blade-like.




Fig. 2. Placoderm and chondrichthyan microremains from the Buchan Group. Scale bar = 100  $\mu$ m unless indicated otherwise. A, *Murrindalaspis* sp. Fragment AMF101198 from Me41.5–42.7m, Pyramids Member of Taravale Formation, *dehiscens* or *perbonus* Zone. B–D, *Buchanosteus* sp. B, Scale AMF101199 from SLO51–68m, Taravale Formation, *perbonus* Zone, crown view. C, Fragment AMF112353 from Me41.5–42.7m, Pyramids Member of Taravale Formation, *dehiscens* or *perbonus* Zone. D, Fragment AMF101197 from G.Rd340.5m, Taravale Formation, *inversus* Zone. E–J, *Goodradigbeeon australium* E, F, Scale AMF112355 from SLO213m, Taravale Formation, *perbonus* Zone. E, crown view, F, lateral view. G–H, Scale AMF101196 from SL46.5m, Buchan Caves Limestone, *dehiscens* Zone. G, crown view, H, laterobasal view. I, J, Scale AMF112356 from SL46.5m, Buchan Caves Limestone, *dehiscens* Zone. I, lateral view, J, crown view. K, *Ohiolepis* sp. Scale AMF101180 from Me91.4–93m, Murrindal Limestone, *perbonus* Zone, crown view. Some of these specimens were illustrated by Basden et al. (2000): A, fig. 10.3; B, fig. 10.4; D, fig. 10.2; G, fig. 10.1; K, fig. 8.9.

Burrow & Turner (1988:687, figs 10B, C) described and illustrated histological features of *Murrindalaspis wallacei* scales.

Order *Arthrodira* Woodward 1891  
Suborder *Brachythoraci* Gross 1932  
Family *Buchanosteidae* White 1952

Genus *Buchanosteus* Stensiö 1945  
*Buchanosteus* sp.  
Fig. 2B–D

*Material.* AMF101199 from SLO51–68m (Taravale Formation, *perbonus* Zone), AMF112353 from Me41.5–42.7m (Pyramids Member of Taravale



Formation, *dehiscens* or *perbonus* zone), AMF101197 from GRd340.5m (Taravale Formation, *inversus* Zone), plus 98 other scales and fragments from SL 39m, 40.8m, 42m (Buchan Caves Limestone, *pirenae* or *dehiscens* Zone), SL 46.5m, 49.7m, 54m, 56.2m, 58.5m (Buchan Caves Limestone, *dehiscens* Zone), SLO 17m, 34m, 51m (Taravale Formation, *dehiscens* Zone), SLO 64m, 51-68m, 213m and G.Rd 51.2m, 51.7m, 52.5m (Taravale Formation, *perbonus* Zone), Mc 4.9-6.2m, 6.5-7.6m, 11-12.2m, 12.5-137m, 14-15.2m, 15.5-18.3m, 17.7m, 18.6-21.3m, 21.6-24.4m, 27.7-30.5m, 35.4-36.6m, 38.4-39.9m, 41.5-42.7m (Pyramids Member of Taravale Formation, *dehiscens* or *perbonus* Zone), Mc 88.7m, 91.4-93m, 144.8-146.3m, 156.1m, 161.5m, 189.9m, 192.5m (Murrindal Limestone, *perbonus* Zone), and G.Rd 340.5m, 343.5m (Taravale Formation, *inversus* Zone).

**Remarks.** The genus has undergone a series of name changes since the holotype was collected near Buchan early last century (Young 1979; Long 1984a, 1995): Chapman (1916) described the specimen as a new species of *Phlyctaenaspis*, it was reassigned by Hills (1936) to *Coccosteus*, and then by Stensiö (1945) to the new genus *Buchanosteus*. White (1952) gave Stensiö's genus a formal diagnosis, and described a similar skull fragment from Burrinjuck as *B. murrumbidgeensis*. White & Toombs (1972) subsequently referred the Burrinjuck material to a new genus, *Parabuchanosteus*. Following study of new specimens from Burrinjuck and re-examination of Chapman's type material, Young (1979) concluded that only one species, *Buchanosteus confertituberculatus* (Chapman), was represented.

In early reports, placoderm scales from Burrinjuck (Ørvig 1969; Giffin 1980) were identified as *Ohioaspis* because of their resemblance to the North American type material of *Ohioaspis tumulosus* Wells 1944. Turner & Murphy (1988) distinguished between *Ohioaspis* scales and similar scales from Nevada, Australia (Burrinjuck) and south China that they called 'buchanosteid-type', because of differences in tubercle morphology.

Burrow & Turner (1998:678-687, figs 2, 3, 4A, B, 5A-J, 6, 9C-G)) assigned scales from the McLarty member of the Murrindal Limestone, collected with dermal plates of *B. confertituberculatus* near Buchan (Long 1991), to the same species as the plates. However, Young et al. (2001) have recently reinterpreted the specimen assigned by Long to *Buchanosteus* as a primitive coccosteid rather than a buchanosteid. Burrow & Turner (1998) recognised

slight morphological and histological differences from the Buchan material in two types of similar scales from Burrinjuck, which were described as Buchanosteidae indet., with tubercles on the crown either rounded and discrete or stellate and tightly-packed. Scales with stellate tubercles, from Longmenshan section, Sichuan Province, China (*dehiscens/perbonus* zones), with similar histology to buchanosteid scales from Burrinjuck, have been described as Buchanosteidae? indet.; it is likely that several primitive placoderm groups possessed scales with stellate tubercles (Burrow et al. 2000). Plate fragments and both types of scales described below resemble microremains of *Buchanosteus confertituberculatus* held in the Australian National University Collections (e.g. sample CPC16965 from an unknown locality near Taemas).

**Description.** Thin, subcircular or subelliptical scales, from 0.7mm diameter to 1.8mm long. Three to 20 tubercles per scale are either discrete and rounded or closely-packed and stellate. Radiating ridges sometimes extend to the tip of the tubercle (Fig. 2B). In some scales the tubercles extend to the margin, although most scales have a thin margin with radiating mucus grooves, producing a denticulate outline (Fig. 2B). The base is gently convex, flat, or gently concave, often with a few small canal openings scattered irregularly over the surface.

Plate fragments (Fig. 2C, D) are ornamented with discrete, rounded, often widely-spaced tubercles, with ridges that can radiate from a rounded top or be subparallel (e.g. some tubercles on AMF101197, Fig. 2B). The bone between the tubercles bears canal openings, especially around the base of the tubercles.

Burrow & Turner (1998, figs 3, 4A, B, 6F, G) illustrated histological features of buchanosteid indet and *Buchanosteus confertituberculatus* scales.

**Genus *Goodradigbeeon* White 1978**  
***Goodradigbeeon australianum* White 1978**  
 Fig. 2E-J

*Goodradigbeeon australianum* White 1978: 175, pl. 6d-f.

*Goodradigbeeon australianum* – Burrow & Turner 1998: 678-80, figs 7, 8A-D.

*Goodradigbeeon* sp. – Basden et al. 2000: 215, 217, figs 10.1, 11.4.

**Material.** AMF101196 and AMF112356 from SL46.5m (Buchan Caves Limestone, *dehiscens* Zone),

AMF12355 from SLO213m (Taravale Formation, *perbonus* Zone) and 30 other scales from SL39m (Buchan Caves Limestone, *pireneae* or *dehiscens* Zone), SL 46.5m, 48.7m, 56.2m, 58.5m (Buchan Caves Limestone, *dehiscens* Zone), SLO8.5m (Taravale Formation, *dehiscens* Zone), SLO213m and GRd51.2m (Taravale Formation, *perbonus* Zone).

**Diagnosis.** Description of ornament by White (1978): Ornamentation ... is tubercular, the tubercles being of the usual lemon-squeezer type with a relatively small enamel cap. When fresh the vertical ridges are seen to be pustular, the pustules at the apex forming a rosette around the cap, and the ridges of the basal intertubercular network are noticeably sharp ... There is no very marked variation in size or shape of the tubercles ... The distribution of the tubercles is close and even ... The scales are covered with minute editions of the tubercles of the general ornament, but there is usually a smooth margin around the surface. Revised description (Burrow & Turner 1998): Scales have between seven and 21 crown tubercles (average 12); the tubercles are from 0.01 to 0.02 mm wide, and bear from seven to ten radial ridges, which often branch at or near the base. The tubercles are separate, with no over-growth evident. Unlike other scales from placoderms, these have small, centrally-positioned tubercles surrounded by larger tubercles. Only rare scales have (weakly developed) mucus grooves peripherally; width of the tubercle-free crown margin varies from negligible, to about one-third of the scale width.

**Remarks.** White (1978) erected the taxon based on macro and microremains collected at localities near Taemas and Wee Jasper in southern New South Wales. Burrow & Turner's (1998) examination of scales from the residue of the holotype held in the collection of the Natural History Museum, London, resulted in an expanded description of the scales, and explanation of the histological differences between scales of this genus and those of *Buchanosteus*.

**Description.** Biconvex scales are subcircular, diamond-shaped or polygonal shaped, with the crown ornamented with 19-34 closely-spaced stellate tubercles. Central tubercles are often smaller. Between seven and ten radiating ridges on tubercles can branch near the base. A tubercle-free zone on the crown margin varies in width from negligible to approximately one-eighth the width of the scale (Fig. 2G). The base has up to three pore openings (Fig. 2H).

Burrow & Turner (1998, figs 4C, 8E) illustrated the histology of *Goodradigbeeon australiamm* scales.

Class **Chondrichthyes** Huxley 1880  
Subclass **Elasmobranchii** Bonaparte 1838  
Order incertae sedis  
**Ohiolepis** Wells 1944

**Diagnosis** (Wells 1944). Small to relatively large shagreen denticles, rounded rhombic, quadrangular, or irregularly elongate or transversely oval in outline, consisting of a broad crown ornamented by numerous spinelike tubercles with enamelled anteriorly indented, anteriorly sloping faces and acute, posteriorly inclined tips, resting directly on an often internally convex, lamellar base....anterior sides or edges of denticles devoid of tubercles over a narrow strip where overlapped by preceding denticle.

**Ohiolepis** sp.  
Fig. 2K

**Material.** AMF101180 from Mc91.4-93m (Murrindal Limestone, *perbonus* Zone), plus three other scales from Mc18.6-21.3m, 35.4-36.6m (Pyramids Member of Taravale Formation, *dehiscens* or *perbonus* Zone), and Mc 91.4-93m (Murrindal Limestone, *perbonus* Zone).

**Remarks.** The genus, established by Wells (1944) from material from USA, has been reported from Australia (Schultze 1968; Giffin 1980; Turner 1982, 1993; Pickett et al. 1985; Young 1993; De Pomeroy 1994; Burrow 1997), China (Wang 1984; Zhu 2000; Zhu et al. 2000) and Germany (Örvg 1969; Friman 1983; Vieth-Schreiner 1983). De Pomeroy (1994) discussed the differences between some of these reported occurrences, in particular the morphology and spacing of the spines on the crown, and the shape and structure of the base.

**Description.** Diamond-shaped scales are 0.5-1.2mm long and 0.5-0.9mm wide. The crown is fairly flat and bears 4-22 closely-packed, conical spines with rounded or pointed tips, that can either cover the crown or leave a smooth, unornamented anterior or anterolateral margin. The anterior face of the spines can be flattened or bear a longitudinal groove that is more developed on the spines with a pointed tip. The groove can extend for about half the length of the spine or almost to the



tip, with spines on the same scale showing variety of groove length. The grooved, pointed spines tend to be more steeply inclined than the smoother, round-tipped spines.

The diamond-shaped base is generally very slightly convex and centrally vaulted. On some specimens, the lateral corners of the base curve down slightly, while on others the base is gently concave with a thin marginal rim. The neck, when present, is shallow and not indented.

Class *Acanthodii* Owen 1846  
Order *Climatiiformes* Berg 1940  
Family *Climatiidae* Berg 1940  
Genus *Nostolepis* Pander 1856  
“*Nostolepis*” *guangxiensis* Wang 1992  
Fig. 3A-D

*Gomphonchus guangxiensis* (Wang) – Turner et al. 1995: 383.

*Gomphonchus?* *Guangxiensis* (Wang 1992) – Burrow 1997: 64-66, pl. 1 figs 4-8, pl. 4 figs 1-6.

“*Nostolepis*” *guangxiensis* Wang 1992 – Burrow et al. 2000: 398-399, pl. 5.4-5.8.

?acanthodian E – Basden et al. 2000: fig. 9.13.

?acanthodian H – Basden et al. 2000: fig. 12.3.

“*Nostolepis*” *guangxiensis* Wang 1992 – Burrow 2002: 91-95, figs 4E-J, 5J-M, 12A, C, D.

**Material.** AMF101193 from Me41.5m (Pyramids Member of Taravale Formation, *dehiscens* or *perbomis* Zone), AMF101190 from Me148.1m (Murrindal Limestone, *perbomis* Zone), plus 36 other specimens from SL46.5m, 58.5m (Buchan Caves Limestone, *dehiscens* Zone), SLO8.5m, 34m (Taravale Formation, *dehiscens* Zone), SLO179m and GRd51.2m (Taravale Formation, *perbomis* Zone), Me7.9m, 15.5m, 21.6m, 27.7m, 38.4m, 41.5m (Pyramids Member of Taravale Formation, *dehiscens* or *perbomis* Zone), Me148.1m, 155.4m, 156.1m (Murrindal Limestone, *perbomis* Zone), and GRd340.5m (Taravale Formation, *inversus* Zone).

**Diagnosis** (Wang 1992; emended by Burrow et al. 2000). Crown flat and rhombic, with strongly developed side ridges converging posteriorly and delineating the upper crown which bears ridges on the anterior part. Sharp ridges also lead forward from the posterior point of the crown, ornamenting the dentinous region of the upper neck/lateral crown. The crown is formed of mesodentine comprising rare cell spaces and


a network of dentine tubules with a well-developed system of fine, vascular canals; only very rare scales have ‘Stranggewebe’, and then only in the lower crown. The scale crown is smaller than the base, and the scale neck is low, clearly marked off from the base. Base is convex, with a rhombic outline, bone-cell lacunae, and radiating Sharpey’s fibres.

**Remarks.** This taxon was erected by Wang (1992) for scales from the Ertang Formation of central Guangxi, southern China (*perbomis* Zone), based partly on his interpretation of mesodentine and Stranggewebe in the crown. Burrow (1997) noted the resemblance of similar scales from several central western New South Wales localities (Trundle Group, *kindlei-dehiscens* zones) to scales from Arctic Canada, assigned by Vieth (1980) to *Gomphonchus* sp. Cf. *G. hoppei*. She also noted that Wang’s (1992) description included plates showing histological features more typical of *Gomphonchus* than *Nostolepis*: dentine tubules extending into the base, a small embryonic growth zone, numerous thin growth zones in the crown, and absence of Stranggewebe in the crown. On these grounds Burrow (1997) tentatively reassigned the species to “*Gomphonchus?*”, and a preliminary report on similar scales from Longmenshan, Sichuan Province, China (*dehiscens-perbomis* zones; Turner et al. 1995) followed this designation.

Meanwhile, Valiukevicius (1994) had reassigned Vieth’s (1980) *Gomphonchus* sp. cf. *G. hoppei* scales to a new species, *Nostolepis laticristata*, a form lacking the Stranggewebe and wide vascular canals in the crown that are usually present in *Nostolepis*. Further examination of the specimens assigned to *G.?* *guangxiensis* specimens from NSW revealed one scale has Stranggewebe, and another has wide vascular canals, thus leading Burrow (in Burrow et al. 2000 and Basden et al. 2000) to revert tentatively to the original nomenclature for the taxon, “*Nostolepis*” *guangxiensis*.

In addition to the scales from China and New South Wales mentioned above, the taxon has been reported from the Point Hibbs Formation, Tasmania (*sulcatus* Zone; Burrow et al. 1998).

**Description. Morphology.** Scales are small, 0.5-0.7mm, usually of similar width and length. The subrhombic crown is smaller than the base (Fig. 3A, C) and either horizontal or sloping upwards towards the posterior. The main diamond-shaped section of the crown is defined by two ridges converging at a posterior





point, and bears up to three or four anterior ribs that sometimes bifurcate at the anterior margin and generally extend almost to the posterior point of the crown. The posterolateral margins of the crown are ornamented by vertical ribs that may also bifurcate near the margin with the neck. The neck is generally shallow and of similar depth anteriorly and posteriorly (Fig. 3B). The neck-base margin is distinctly marked by a thin rim. The moderately to deeply convex base is centrally vaulted.

**Histology.** The base, a low-angled cone, contains bone cell lacunae and radiating Sharpey's fibre bundles. Growth zones are clearly visible in both base and crown. The crown contains Strangewebe and a network of fine branching dentine tubules.

cf. *Nostolepis* sp.

Fig. 3E, F

**Material.** AMF101192 from SLO34m (Taravale Formation, *dehiscens* Zone), plus 13 other specimens from G.Rd52.5m (Taravale Formation, *perbonus* Zone), and Me3m, 4.9m, 6.5m, 7.9m, 14m, 27.7m, 33.8m (Pyramids Member of Taravale Formation, *dehiscens* or *perbonus* Zone).

**Remarks.** See Denison (1979) for diagnosis and De Pomeroy (1996) for comments on the distribution of this cosmopolitan genus. *Nostolepis* scales have been recorded from the Early Devonian of eastern Australia in the Broken River Group (*serotinus* Zone) and Martins Well Limestone (*pesavis-sulcatus* zones) in Queensland (Turner 1991, 1993; De Pomeroy 1996; Turner et al. 2000), and the Condobolin Formation, Garra and Windellama limestones, Trundle and Yarra Yarra Creek groups in New South Wales, in horizons dated from Lochkovian to early Emsian (Turner 1991, 1993; Burrow 1997; Basden et al. 2000).

**Description.** Small scales 0.4–0.7mm long and wide. The subtriangular crown slopes up from a rounded anterior margin to a pointed posterior that extends beyond the base (Fig. 3E). Short, low, parallel ridges ornament the anterior part of the crown. The neck is shallow anteriorly and deeper and concave posteriorly (Fig. 3F), with no or few visible pore canal openings. The base and neck are separated by a distinct rim (Fig. 3F). The centrally convex base is subrhombic, with anterior and posterior corners slightly rounded and lateral corners more pointed (Fig. 3E). Faint concentric lines visible on the base possibly represent different growth zones.

Genus *Nostolepoides* Burrow 1997

*Nostolepoides platymarginata* Burrow 1997

Fig. 3G–L

*nostolepid* scale – Philip 1965, pl. 8.38. – Turner 1991: pl. 3B.

*acanthodian* scale – Turner 1991: pl. 5H.

*Nostolepoides platymarginata* Burrow 1997: 45–57, pl. 1 figs 1–3, pl. 2 figs 7–10, 12.

*Nostolepoides platymarginata* Burrow 1997 – Basden 1999b: 535, pl. 2 figs 2–3.

*Nostolepoides platymarginata* Burrow 1997 – Burrow 2002: 83–87, figs 5D–I, 8A–K.

**Material.** AMF101183 from SLO8.5m, AMF112378 from SLO51m (Taravale Formation, *dehiscens* Zone), AMF101185 from SLOc85m (Taravale Formation, *perbonus* Zone), plus 396 specimens from SL39m, 39.7m, 40.8m (Buchan Caves Limestone, *pirenae* or *dehiscens* Zone), SL 46m, 48.7m, 58.55m, (Buchan Caves Limestone, *dehiscens* Zone), SLO8.5m, 17m, 34m, 44m, 51m (Taravale Formation, *dehiscens* Zone), SLO55m, 64m, 68m, 72m, 85m, 170m, 179m, 213m and G.Rd 51.2m, 52.5m (Taravale Formation, *perbonus* Zone), Me3.4m, 4.9m, 7.9m, 11m, 15.5m, 18.6m, 21.6m, 24.2m, 27.7m, 30.8m, 36.9m, 41.5m,

Fig. 3. Acanthodian microremains from the Buchan Group. Scale bar = 100µm. A–D, “*Nostolepis*” *guangxiensis*. A, B, Scale AMF101193 from Me41.5–42.7m, Pyramids Member of Taravale Formation, *dehiscens* or *perbonus* Zone. A, crown view, B, lateral view. C, D, Scale AMF101190 from Mc148.1–149.4m, Murrindal Limestone, *perbonus* Zone. C, crown view, D, lateral view. E, F, cf. *Nostolepis* sp. Scale AMF101192 from SLO34m, Taravale Formation, *dehiscens* Zone. E, crown view, F, lateral view. G–L, *Nostolepoides platymarginata*. G, H, Scale AMF101183 from SLO8.5m, Taravale Formation, *dehiscens* Zone. G, crown view, H, lateral view. I, J, Scale AMF112378 from SLO51m, Taravale Formation, *dehiscens* Zone. I, crown view, J, lateral view. K, L, Scale AMF101185 from SLO85m, Taravale Formation, *perbonus* Zone. K, crown view, L, lateral view. M, *Cheiracanthoides* sp. cf. *C. comptus*. Scale AMF101215 from Me88.7m, Murrindal Limestone, *perbonus* Zone, crown view. N, O, *Cheiracanthoides* sp. cf. *C. wangi*. N, Scale AMF101214 from SLO213m, Taravale Formation, *perbonus* Zone, crown view. O, Scale AMF101216 from Mc160.6m, Murrindal Limestone, *perbonus* Zone, crown view. P, Q, *Rhadinacanthus*? sp. Scale AMF101189 from SL46.5m, Buchan Caves Limestone, *dehiscens* Zone. P, crown view, Q, lateral view. R, *Gomphonchus bogongensis*. Scale AMF101217 from SLO17m, Taravale Formation, *dehiscens* Zone, crown view. Some of these specimens were illustrated by Basden et al. (2000): A, fig. 9.11; C, fig. 9.8; E, fig. 9.10; G, fig. 9.1; K, fig. 9.3; P, fig. 9.7.

43m (Pyramids Member of Taravale Formation, *dehiscens* or *perbonus* Zone), Mc88.7m, 91.4m, 150.9m, 154m, 155.4m, 156.1m, 156.4m, 160m, 160.6m, 163.8m, 180.1m, 182.9m, 189.9m, 190.5m, 192.5m, 227.4m (Murrindal Limestone, *perbonus* Zone), GRd 340.5m (Taravale Formation, *inversus* Zone).

**Diagnosis** (Burrow 1997). Scales range from 0.15 to 1.0mm wide. They have a bi-level crown; the upper crown is delineated by ridges converging in the posterior half of the crown and with subparallel ribs between them. The lower crown is a latero-posterior, overhanging ledge of varying width and outline. The neck is high and concave (morphotype 1) or low anteriorly (morphotype 2), and the base is moderately vaulted. The crown is composed of mesodentine. Large bone cell lacunae and their processes interconnect throughout the base.

**Remarks.** These Victorian scales resemble Morphotype 1 scales described by Burrow (1997) from the Gleninga (*pireneae-dehiscens* zones) and Troffs formations (*kindlei-dehiscens* zones) in central NSW. They are morphologically similar to the Late Silurian-Early Devonian *Nostolepis striata* Pander but can be distinguished from that species histologically and by the posterolateral ledge of the crown being wide and/or irregular and continuing around the posterior point, as discussed by Burrow (1997). One of the scales figured by Philip (1965, pl. 8.38) from the Tyers area has been assigned by Burrow (1997) to *Nostolepoides platymarginata*.

**Description. Morphology.** Scales range in width from 0.5 to 1.2mm. The subrhombic crown consist of two levels. The higher central triangular section is enclosed by two strong radial ridges extending from the anterior margin and converging in a posterior point (Fig. 3G, I), with 3-5 short anterior subparallel ribs that extend up to halfway towards the posterior point, and can bifurcate at the anterior margin (Fig. 3I). The lower level of the crown is a thin posterolateral platform of varying width, that extends beyond the posterior point of the upper crown and can have an irregular (Fig. 3I) or even denticulate margin (Fig. 3G, K). The fairly deep neck is clearly defined, concave, and deeper posteriorly. Some scales have a row of up to ten pore canal openings on the anterior neck (Fig. 3H). The subrhombic base is gently convex, centrally vaulted and separated from the neck by a distinct rim (Fig. 3H, J, L).

**Histology.** Thin sections show a clear demarcation between crown and base, with a large initial (embryonic) growth zone. Both base and crown have distinct growth zones. The base contains bone cell lacunae and radiating bundles of Sharpey's fibres. The branching dentine tubules in the crown are confined to individual growth zones and are less apparent in outermost zone(s).

Genus **Cheiracanthoides** Wells 1944

**Cheiracanthoides** sp. cf. *C. comptus*

Fig. 3M

**Material.** AMF101215 from Mc88.7m (Murrindal Limestone, *perbonus* Zone), and 93 other scales from SL39m, 40.8m (Buchan Caves Limestone, *pireneae* or *dehiscens* Zone), SL46.5m, 58.6m (Buchan Caves Limestone, *dehiscens* Zone), SLO 8.5m, 17m, 34m, 51m (Taravale Formation, *dehiscens* Zone), SLO 55m, 64m, 72m, 85m, 213m and GRd50.9m, 52.8m (Taravale Formation, *perbonus* Zone), Mc 15.5m, 33.8m (Pyramids Member of Taravale Formation, *dehiscens* or *perbonus* Zone), Mc88.7m, 156.1m, 160.6m, 189.9m, 190.5m (Murrindal Limestone, *perbonus* Zone), GRd339.7m (Taravale Formation, *inversus* Zone).

**Remarks.** The cosmopolitan species *Cheiracanthoides comptus* is frequently reported from Early and Middle Devonian horizons (e.g. Wells 1944; Gross 1973; Denison 1979; Valiukevicius 1979, 1985, 1994, 1998; Vieth 1980; Vieth-Schreiner 1983; Poltnig 1984; Boucot et al. 1989; Turner 1991, 1993; Wang 1992; De Pomcroy 1995, 1996; Basden et al. 2000; Burrow et al. 2000; Zhu 2000; Zhu et al. 2000). Scales from the *Receptaculites* Limestone at Burrinjuck have been assigned to the genus (Giffin 1980, fig. 5), but it is possible that *C. comptus* has become a "bucket" taxon comprising several different form species or even genera with a similar morphology; see De Pomcroy (1996) for discussion.

Recent work has resulted in the erection of several new species (Valiukevicius 1998, Burrow et al. 2000). The diagnosis and description, originally established by Wells (1944), have been refined so that more subtle differences in morphology and histology can be attributed to different species. For example, from material from the East Baltic and Byelorussia, Valiukevicius (1998) has proposed three new species – *Cheiracanthoides planis* (smaller than *C. comptus*, with fewer ridges on the crown and absence of the



narrow unsculptured rim around the crown anterior), *C. nativus* (lacking the smooth lateral areas on the crown of *C. comptus*) and *C. estonicus* (scales from the Eastern Baltic previously assigned to *C. comptus*). Burrow et al. (2000), studying residues from Sichuan Province, China, have distinguished *C. dolosus*, with a bevelled anterior crown rim, short regularly-spaced ridges, and Stranggewebe extending into the anterior crown, and *C. wangi* (discussed below).

**Description.** The elliptical or subrhombic crown is flat, slightly wider than long, and ornamented on the anterior third to half by between four and ten radiating or subparallel low rounded ridges separated by broad furrows. The indented neck is a consistent depth from anterior to posterior. The crown extends beyond the base posteriorly. The base is diamond-shaped, moderately convex and centrally vaulted, with the neck-base margin marked by a thickened rim.

**Cheiracanthoides wangi** Burrow et al. 2000

**Cheiracanthoides** sp. cf. *C. wangi*

Fig. 3N, O

**Material.** AMF101214 from SLO213m (Taravale Formation, *perbonus* Zone), AMF101216 from Me160.6m, and two scales from Me88.7m, 160.6m (Murrindal Limestone, *perbonus* Zone).

**Diagnosis.** Scales with a fairly flat crown, which is more than twice as long as wide, and with anterior edges often forming a sharp angle. The anterior crown is ornamented with between three and ten sharp, strongly-developed, sub-convergent ridges which extend about one-third of the crown length. The lateral areas of the anterior crown are smooth: i.e., the ridges span the medial two-thirds to three-quarters of the anterior crown. More than half of the crown overhangs the posterior limit of the base; the overhanging section is very thin. The neck is concave all round, falling away sharply from the anterior crown edge, and deepening slightly towards the posterior. The base/neck junction is a strongly marked rim; the base is moderately to strongly vaulted, and projects slightly forwards of the crown. "Stranggewebe" is only formed in the upper neck/crown area of the posterior growth zones. Bone cell lacunae are preserved in the apex of the base (Burrow et al 2000).

**Remarks.** The species was described by Burrow et al. (2000: 397-398, figs 2.10-2.12, pl. 3.1-3.7, 3.9-

3.10) from Longmenshan, Sichuan Province, China (*dehiscens-perbonus* conodont zones), and is also reported from the late Pragian – early Emsian (?*kindlei-perbonus* zones) in the Guangxi and Sichuan provinces (Zhu 2000; Zhu et al. 2000). It is now considered an index fossil for the *dehiscens* Zone (Burrow 2000).

The Australian scales described here differ in some features from the Chinese specimens. The central region of the crown generally tends to have ridges closer to parallel, and with a more distinct difference between this central parallel section and the anterior section where the ridges radiate; and the length of the crown, which in some scales is greater than the width, is, even in broken specimens, generally nowhere near twice the width, as is the case with *C. wangi* scales from China. However, since all the Victorian specimens have the thin posterior section of the crown broken off, it is difficult to determine precisely the initial dimensions and length/width ratios of the scales.

**Description. Morphology.** Scales are approximately 0.8 to 1.2 mm wide, and 0.9 to 1.2 mm from anterior margin to broken posterior edge. The flat subcircular or subrhombic crown slopes slightly upwards towards the posterior. The anterior part of the crown bears between five and seven sharply-crested, radiating ridges that become closer together and subparallel in the central crown area. At the anterior, these ridges can extend down a short distance onto the upper part of the neck. The posterior part of the crown is unornamented, thin and commonly broken.

The neck is generally high and concave, and can become deeper posteriorly. Several small round pore canal openings are found in the central part of the anterior neck, and more elongate openings are high up under the overhanging crown on the posterior neck. The base is very deeply convex, and centrally vaulted. The base and neck are separated by a swollen rim that extends into lateral and posterior flanges.

**Histology.** The base inserts high into the crown. Both base and crown have distinct growth zones, those in the base parallel to the lower surface. The base contains bone cell lacunae with branching tubules and radiating Sharpey's fibre bundles. All but the outermost growth zones of the crown contain dense outwardly-directed tubules, all confined within individual growth zones.

Family **Diplacanthidae** Woodward 1891

Genus **Rhadinacanthus** Traquair 1888

**Rhadinacanthus?** sp.

Fig. 3P, Q

**Material.** AMF101189 from SL46.5m (Buchan Caves Limestone, *dehiscens* Zone) and seven specimens from SL39m (Buchan Caves Limestone, *pireneae* or *dehiscens* Zone), SL46.5m (Buchan Caves Limestone, *dehiscens* Zone), SLO51-68m (Taravale Formation, *perbonus* Zone), and Mc160.6m (Murrindal Limestone, *perbonus* Zone).

**Remarks.** The genus *Rhadinacanthus* is considered a possible synonym for *Diplacanthus* (Denison 1979; see Burrow et al. 2000, for discussion).

**Description.** Scales have a kite-shaped crown that is wider than long. The flat crown has a curved anterior margin and pointed lateral and posterior corners (Fig. 3P). The posterior crown overhangs the base only a short distance (Fig. 3Q). Approximately 15 closely-spaced sharp-crested anterior ribs on the crown converge slightly towards the crown centre and extend back to about mid-way.

The indented neck is clearly defined from both crown and base by a distinct rim. The neck has up to ten pore openings low on the neck close to the neck-base margin both anteriorly and posteriorly, but shows no sign of the wart-like bulges at the posterior, as described by Gross (1973) in *R. balticus*. The base is gently convex.

Order *Ischnacanthiiformes* Berg 1940  
Family *Ischnacanthidae* Woodward 1891  
Genus *Gomphonchus* Gross 1971  
*Gomphonchus? bogongensis* Burrow 1997  
Fig. 3R

*Gomphonchus? bogongensis* Burrow 1997: 67-71, pl. 1 figs 9-12, 16, pl. 4 figs 8-16.

*Gomphonchus? bogongensis* Burrow 1997 – Burrow et al. 1998: 13, fig. 3B-C.

*Gomphonchus? bogongensis* Burrow 1997 – Basden 1999b: 536-8, pl. 2 figs 8-9.

*Ischnacanthid* indet. C – Lindley 2002: fig. 9A-I.

**Material.** AMF101217 from SLO17m (Taravale Formation, *dehiscens* Zone), and 152 specimens from SLO39m, 40.8m (Buchan Caves Limestone, *pireneae* or *dehiscens* Zone), SL46.5m, 48.7m, 53.2m, 54m, 56.2m, 58.5m (Buchan Caves Limestone, *dehiscens* Zone), SLO8.5m (Taravale Formation, *dehiscens* Zone), SLO170m, 213m, G.Rd51.7m, 52.5m

(Taravale Formation, *perbonus* Zone), Mc 4.9m, 6.5m, 8.5m, 11m, 14m, 15.5m, 18.6m, 30.8m, 33.8m, 35.4m, 36.9m, 38.4m, 43m, 44.5m (Pyramids Member of Taravale Formation, *dehiscens* or *perbonus* Zone), Mc150.9m, 154m, 155.4m, 156.1m (Murrindal Limestone, *perbonus* Zone).

**Diagnosis** (Burrow 1997). Small scales, 0.15 to 0.7mm wide, with a subtriangular crown, low neck, and a moderate to large convex base. There are two main morphotypes: in morphotype 1, the crown has a curved fan shape, ornamented with sharp fan-like ridges and grooves, and slopes upwards from the anterior edge to a posterior point. Morphotype 2 scales have a crown with an upper sub-triangular, and a lower lateral, level which can be greatly elongated posteriorly. The scale crown is composed of orthodentine. All scales have a 'scalloped' antero-lateral crown edge.

**Remarks.** Scales resemble Morphotypes 1 and 2 described by Burrow (1997) from the Trundle Beds (now Troffs Formation) and Gleninga Formation in central NSW. Revised dating of the horizons by Sherwin (1996) now places the horizons with these scales into *pireneae-dehiscens* zones. Scales of *G? bogongensis* are also reported from the *sulcatus* Zone at Point Hibbs (Burrow et al. 1998).

Morphotypes 1 and 2 can both be distinguished from *Nostolepoides platymarginata* scales by the scalloped anterior margin of the crown, and the increased depth of the neck towards the posterior, and Morphotype 1 also by the lack of lowered lateral crown ledges. Morphotype 1 is distinguished from "*Nostolepis*" *guangxiensis* by the short anterolateral ridges not continuing around below the posterior point of the crown (Burrow 1997).

**Description. Morphology.** Subrhombic or diamond-shaped scales have the crown and base approximately a similar shape, and the crown generally smaller than the base. Width ranges from 0.4 to 1.2mm. The fan-shaped crown has 4-8 radiating anterior ridges with sharp crests, separated by deep furrows; these ridges can be less than half the crown length, or extend virtually to the posterior point of the crown. These features in combination with the curving down of the anterior crown towards the neck give the scalloped anterior margin to the crown that is distinctive of this form.

In Morphotype 2 scales the posterior point of the crown is more elongated than in Morphotype 1, and can extend slightly beyond the base. Short anterolateral



ridges on the sides of the crown do not continue beyond the posterior point. These anterolateral ridges vary, from very short anterior ridges that give a bifurcating appearance to the marginal lateral ribs of the crown, or do not even quite meet the marginal lateral ribs, to ridges found high on the lateral margins of the crown, not extending far onto the neck.

The neck is short anteriorly, not very indented, and widens posteriorly. A row of pore canal openings on the posterior neck is better developed in some scales, with the canal openings elongated.

The base is moderately to strongly convex, centrally vaulted and separated from the neck by a distinct rim that is particularly pronounced at the lateral and posterior corners. Concentric markings parallel to the neck/base rim indicate presence of Sharpey's fibres.

**Histology.** Scales are generally not well preserved, with minimal histological detail revealed in many scales. However, growth zones in both base and crown are apparent. The base contains Sharpey's fibre bundles, a fine network of interconnecting tubules, and sparse bone cell lacunae. The crown contains fine branching tubules that are sparse in the outermost growth zone.

Class *Osteichthyes* Huxley 1880  
Subclass *Aetinopterygii* Klein 1885  
Order *Palaeonisciformes* Hay 1929  
  *Ligulalepis* Schultz 1968  
  *Ligulalepis toombsi* Schultz 1968  
    Fig. 4A

*Ligulalepis toombsi* Schultz 1968: 345-351, figs 1-6, pl. 1 figs 1-8, pl. 4, fig. 1.

*Ligulalepis toombsi* Schultz 1968 – Giffin 1980: 10-11, fig. 10.

*Ligulalepis toombsi* Schultz 1968 – Burrow 1994: 177-184, figs 2-5.

*Ligulalepis toombsi* Schultz 1968 – Basden et al. 2000: fig. 8.7.

**Material.** AMF 101178 from SLO34m (Taravale Formation, *dehiseens* Zonc).

**Diagnosis** (translated from Schultz 1968). Genus ... with deep scales on the flanks; scales close to the shoulder girdle are 3-4 times deeper than long; on the anterior upper corner there is a very marked projection, which makes a sharp angle to the front edge; ventral edge slopes away from the front edge towards the back

end. Strong peg and correspondingly deep socket; keel quite deep; secondary keel, on the back edge of which are openings of most canals. Development of the ganoin surface common to all species in the genus. Ganoin well-developed, partly divided into separate ribs backwards from front edge, ornamented towards front edge with small longitudinal striations which are absent in small scales; back edge comb-like; front edge slightly or strongly concave, anterior lower corner rounded, ventral edge straight.

**Remarks.** Schultz (1968) erected the taxon from material in residues from the 1955 and 1963 Burrinjuck expeditions by H.A. Toombs (Natural History Museum, London), from spot localities in horizons from the Cavan Formation to the Warroo Limestone. Giffin (1980) figured similar scales from the *Receptaculites* Limestone at Taemas (residues of B.D.E. Chatterton's processing of trilobites and brachiopods) as did Burrow (1994) from the Troffs and Gleninga formations of central New South Wales (*pireneae-dehiseens* zones). Other Devonian palaeoniscoid genera described from Australia are *Terenolepis turnerae* Burrow 1995 from Connemarra Formation, Trundle Group and Garra Limestone (Hocking in Basden et al 2000), *Howqualepis rostridens* from the Frasnian Avon River Group at Mt Howitt, Victoria (Long 1988, figs 32, 33), and *Mimia toombsi* and *Moythomasia durgaringa* from the Givetian-Frasnian Gneudna Formation and Frasnian Gogo Formation (Gardiner & Bartram 1977; Trinajstić 1999a, figs 3, 4; 1999b, figs 4, 5).

**Description.** The subrectangular scale has a free field ornamented with unfused ridges, a denticulate caudal margin and a smooth depressed field along the curved rostral edge. A rounded, tongue-shaped process projects rostro-dorsally and more pointed dorsal peg would have fitted into a corresponding socket on the ventral edge of the scale above. On the basal surface dorsoventrally oriented primary and secondary keels are separated by a groove.

Subclass *Sarcopterygii* Romer 1955  
Division *Rhipidistia* Cope 1887  
  Order *Onychodontiformes*  
  *Onychodontiformes* gen. et sp. indet.  
    Fig. 4B-J

**Material.** Scales with *U-shaped tubereles*: AMF118232 from Mc7.9-9.1m (Pyramids Member




Fig. 4. Osteichthyan microremains from the Buchan Group. Scale bar = 100µm unless indicated otherwise. A, *Ligulalepis toombsi* Scale AMF101178 from SLO34m, Taravale Formation, *dehiscens* Zone. B-J, Onychodontiiformes gen. et sp. indet. B, Fragment AMF118232 from Me7.9-9.1m, Pyramids Member of Taravale Formation, *dehiscens* or *perbonus* Zone, crown view. C, Fragment AMF101181 from Me156.1m, Murrindal Limestone, *perbonus* Zone, crown view. D, E, Fragment AMF118231 from GRd340.5m, Taravale Formation, *perbonus* Zone. D, crown view, E, detail of crown ornament. F, Tooth AMF101220 from SLO55m, Taravale Formation, *perbonus* Zone. G, H, Scale AMF101182 from Me43-44.2m, Pyramids Member of Taravale Formation, *dehiscens* or *perbonus* Zone. G, crown view, H, lateral view. I, J, Scale AMF118236 from GRd340.5m, Taravale Formation, *inversus* Zone. I, crown view, J, coronolateral view. Some of these specimens were illustrated by Basden et al. (2000): A, fig. 8.7; C, fig. 8.10; G, fig. 8.11; H, fig. 8.12.



of Taravale Formation, *dehiscens* or *perbonus* Zone), AMF101181 from Me156.1m (Murrindal Limestone, *perbonus* Zone), AMF118231 from G.Rd340.5m (Taravale Formation, *perbonus* Zone), plus 37 other specimens from G.Rd340.5m (Taravale Formation, *perbonus* Zone), Me3m, 7.9-9.1m (Pyramids Member of Taravale Formation, *dehiscens* or *perbonus* Zone), Me151.5m (Murrindal Limestone, *perbonus* Zone). *Teeth*: AMF101229 from SLO55m (Taravale formation, *perbonus* Zone), plus 157 other specimens from SLO8.5m, 17m, 34m, 44m, 51m (Taravale Formation, *dehiscens* Zone), SLO64m, 213m, G.Rd50.9m, 52.3m (Taravale Formation, *perbonus* Zone), Me3m, 3.4-3.6m, 4.9-6.2m, 6.5-7.6m, 7.9-9.1m, 11-12.2m, 12.5-13.7m, 14-15.2m, 15.5-18.3m, 17.7m, 18.6-21.3m, 21.6-24.4m, 24.4-27.4m, 27.7-30.5m, 30.8-33.5m, 33.8-35.1m, 35.4-36.6m, 36.9-38.1m, 38.4-39.9m, 43-44.2m (Pyramids Member of Taravale Formation, *dehiscens* or *perbonus* Zone), Me91.4-93m, 150.9m, 151.5m, 155.4m, 156.1m, 156.4m, 160.6m, 161.5m, 163.1m, 163.8m, 192.5m (Murrindal Limestone, *perbonus* Zone), Me224-225.6m, 225.9-227.1m, 227.4-228.6m (Upper Taravale Formation, *perbonus* Zone), G.Rd340.5m (Taravale Formation, *inversus* Zone).

*Scales with curved denticle*: AMF101182 from Me43-44.2m (Pyramids Member of Taravale Formation, *dehiscens* or *perbonus* Zone), AMF118236 from G.Rd340.5m (Taravale Formation, *inversus* Zone).

*Remarks*. Inclined U-shaped tubercles with a central depression are found on scales of several fish groups, thus limiting their utility for identification. Dipnoan scales have been described with this type of tubercular ornament in conjunction with eosmine and Westoll lines (the primitive dipnoan *Dipnorhynchus sussnilehi* from Buchan and Taemas; Thomson & Campbell 1971), but as no eosmine is preserved on the Victorian scales, dipnoan affinity is probably precluded. Apart from dipnoans, the only other common osteichthyan remains known from the region are actinopterygians and onychodontids, and since actinopterygians are not known to have denticles, the scales and fragments described here probably belong to onychodonts.

*Description. Scales with U-shaped tubercles*. Figured specimens are broken fragments that exhibit a variety of morphologies of the crescent- or U-shaped tubercles. The tubercles themselves can be fairly flat with smooth, very slightly concave anterior margins (Fig. 4D),

oriented in rows (Fig. 4C), or more elongate and irregularly shaped with a distinct depression in the anterior face (Fig. 4B).

The detail of specimen AMF118231 (Fig. 4E) shows the bone surface between the dentine tubercles pitted with irregular vascular canal openings, whereas in AMF101181 (Fig. 4C) the bone between the tubercles is striated by rows of elongate canal openings.

*Teeth*. Simple curved conical teeth up to 2mm long can be slender, tapering to a sharp point, or thicker with a rounded apex (Fig. 4F). The base of the lanial and interlanial teeth is circular and constricted. All specimens are hollow, with the cusp in better-preserved examples covered in finely striated enamel, although preservation is not sufficient to discern micro-patterns in the enamel.

*Scales with curved denticle*. Articulated diamond-shaped scales (Fig. 4G) have a central curved, inclined, conical, unstriated denticle. The main part of the scale consists of flat bony tissue; in contrast, the denticles are noticeably smoother and shinier. The only ornament on the scales is a series of faint radiating grooves on the base around each denticle (Fig. 4I, J). The basal surface of the scales generally is flat and unornamented, although specimen AMF101182 (Fig. 4G, H) has faint grooves corresponding to the margins of the articulated diamond-shaped scales, and the three articulated scales in specimen AMF118236 (Fig. 4I, J) are slightly concave centrally. A broken denticle (Fig. 4I, J) shows a very narrow longitudinal canal, situated slightly off-centre towards the outer edge of the denticle.

## CONCLUDING REMARKS

Acanthodian scales are the most common elements of the fauna (Fig. 3). Climatid scales of *Nostolepoides platymarginata* and *Cheiracanthoides* sp. cf. *C. eomptus* are spread throughout the three sections, and occur in all four formations (Buchan Caves and Murrindal limestones, Taravale Formation, and the Pyramids Member of the Taravale Formation). In SL/SLO section, *Gomphonehus? bogongensis* is much more common in the Buchan Caves Limestone.

The "nostolepid" scale (*sulcatus* Zone) from Tyers in Victoria, illustrated by Philip (1965) and Turner (1991, pl. 3B), is similar to some of these scales, but another similar scale was assigned by Vieth (1980, pl. 6, fig. 15) to *Cheiracanthoides eomptus*. More elongate scales (e.g. Fig. 3E, F) resemble the scale from

Burrinjuck in southern New South Wales assigned to *C. comptus* by Giffin (1980, fig. 5), but these are also similar to a scale from Algeria assigned by Blicek et al. (1984, pl. 1, fig. 4) to *Nostolepis* sp. Similar crown morphologies but different shape (Fig. 3G-L) might indicate scales from different parts of the same fish, but V.T. Young (1995) recorded little shape variation in several articulated acanthodian taxa from the Early to Middle Devonian of Britain. Other scales are compared with *Cheiracanthoides wangi* (Fig. 3N, O), recently described from *dehiscens-perbonus* zones of China (Burrow et al. 2000), and *Rhadinacanthus* (Fig. 3P, Q).

Scales of arthrodires *Goodradigbeeon* and *Buchanosteus* are the most abundant placoderm microremains in the Buchan measured stratigraphic sections. Small plate fragments (Fig. 2E, F) recorded from horizons in the *dehiscens* – *inversus* interval have the same ornament as originally figured by White (1952, pl. 30) for *Buchanosteus murrumbidgeensis*. Other plate fragments with more elongate tubercles (Fig. 2A) are compared with *Murrindalaspis* sp. (e.g. Long & Young 1988).

Osteichthyan remains from Buchan include a scale of the actinopterygian *Ligulalepis toombsi* (Fig. 4A), and cosmine covered elements that may belong to either dipnoans or porolepiforms (see De Pomeroy, 1996: 433). Similar cosmine-covered scales described from an Early Devonian dipnoan by Barwick et al. (1997) have rows of odontodes as seen in porolepiform scales, (e.g. Fig. 4B-E; Ørvig, 1969, fig. 6). However, the scales of *Dipnorhynchus*, the only named dipnoan from Buchan, have a different morphology, with coarser rounded tubercles on the scale margins (see Thomson & Campbell 1971, figs 86-89). By the Late Devonian these tubercle zones had been lost from dipnoan scales (Pridmore & Barwick 1993). Rhombic plates, each with a central hooked denticle (Fig. 4G-J) may be small elements from inside the bucco-pharyngeal cavity of an onychodontid (P. Janvier, pers. comm.). In section SL/SLO, both osteichthyans present (*Ligulalepis toombsi* and *Onychodus* sp.) occurred only in the Taravale Formation, and were absent from the Buchan Caves Limestone.

A chondrichthyan scale similar to *Ohiolepis* sp. (Fig. 2K) is the first chondrichthyan reported from the Buchan sequence.

The first thelodont scales described from the Buchan area were part of the current study (Basden 1999a). All ten scales were recovered from horizons dated *dehiscens-perbonus* zones, and have been

assigned to *Turinia* sp.cf. *T. australiensis*.

## ACKNOWLEDGEMENTS

I thank Professors Ruth Mawson and John Talent for making available the microvertebrate material for study, for providing the research facilities to complete my Ph.D., of which this project formed a part, and for giving ongoing encouragement and assistance. I thank Dr Gavin Young for continual encouragement and advice, Drs Carole Burrow and Susan Turner for helpful discussion, and two anonymous referees for constructive comments on an earlier manuscript. This work was partly carried out while in receipt of an Australian Postgraduate Award.

## REFERENCES

- BASDEN, A.M., 1999a. Emsian (Early Devonian) microvertebrates from the Buchan and Taemas areas of southeastern Australia. *Records of the Western Australian Museum Supplement* 57: 15-21.
- BASDEN, A.M., 1999b. Early Devonian microvertebrates from the Tyers-Boola area of central Victoria, Australia. *Bollettino della Società Paleontologica Italiana* 37(2-3): 527-541.
- BASDEN, A.M., BURROW, C., HOCKING, M., PARKES, R. & YOUNG, G.C., 2000. Siluro-Devonian microvertebrates from southeastern Australia. In *Palaeozoic Vertebrate Biochronology and Global Marine/Non-marine Correlation - Final Report of IGCP 328 (1991-1996)*, Blicek, A. & Turner, S., eds, *Courier Forschungsinstitut Senckenberg* 223: 201-222.
- BLIECK, A., GOUJET, D., JANVIER, P. & LELIÈVRE, H., 1984. Microrestes de vertébrés du Siluro-Dévonien d'Algérie, de Turquie et de Thaïlande. *Geobios* 17: 851-856.
- BOUCOT, A.J., MCCLURE, H.A., ALVAREZ, F., ROSS, J.R.P., TAYLOR, D.W., STRUVE, W., SAVAGE, N.N. & TURNER, S., 1989. New Devonian fish fossils from Saudi Arabia and their biogeographical affinities. *Senckenbergiana lethaea* 69: 535-597.
- BURROW, C.J., 1994. Form and function in scales of *Ligulalepis toombsi* Schultze, a palaeoniscoid



- from the Early Devonian of Australia. *Records of the South Australian Museum* 27: 175-185.
- BURROW, C.J., 1997. Microvertebrate assemblages from the Lower Devonian (*pesavis/sulcatns*) of central New South Wales, Australia. *Modern Geology* 21: 43-80.
- BURROW, C.J., 2000. Changes in microvertebrate faunal associations through the Lower Devonian of south-eastern Australia. *Palaentology Down Under 2000. Geological Society of Australia, Abstracts* 61: 149-150.
- BURROW, C.J. & TURNER, S., 1998. Devonian placoderm scales from Australia. *Journal of Vertebrate Paleontology* 18(4): 677-695.
- BURROW, C.J. & TURNER, S., 1999. A review of placoderm scales, and their significance in placoderm phylogeny. *Journal of Vertebrate Paleontology* 19(2): 204-219.
- BURROW, C.J., LONG, J.A. & TURNER, S., 1998. Lower Devonian microvertebrates from the Point Hibbs Formation, Tasmania. *Alcheringa* 22: 9-20.
- BURROW, C., TURNER, S. & WANG, S.T., 2000. Devonian microvertebrates from Longmenshan, Sichuan, China: taxonomic assessment. In *Palaeozoic Vertebrate Biochronology and Global Marine/Non-marine Correlation - Final Report of IGCP 328 (1991-1996)*, Blicek, A. & Turner, S., eds, *Courier Forschungsinstitut Senckenberg* 223: 391-451.
- CHAPMAN, F., 1916. On the generic position of *Asterolepis ornata* var. McCoy, with description of a new variety. *Proceedings of the Royal Society of Victoria* 28: 211-215.
- DENISON, R.H., 1979. Acanthodii. In *Handbook of Paleichthyology*, Schultz, H.P. (ed), Gustav Fischer Verlag, Stuttgart.
- DE POMEROY, A.M., 1994. Mid-Devonian chondrichthyan scales from the Broken River, North Queensland, Australia. *Memoirs of the Queensland Museum* 37(1): 87-114.
- DE POMEROY, A.M., 1995. Australian Devonian fish biostratigraphy in relation to conodont zonation. *Courier Forschungsinstitut Senckenberg* 182: 475-486.
- DE POMEROY, A.M., 1996. Biostratigraphy of Early and Middle Devonian microvertebrates from Broken River, north Queensland. *Records of the Western Australian Museum* 17: 417-437.
- FRIMAN, L., 1983. *Ohiolepis*-Schuppen aus dem unteren Mitteldevon der Eifel (Rheinisches Schiefergebirge). *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte* 1983(4): 228-236.
- GARDINER, B.G. & BARTRAM, A.W.H., 1977. The homologies of ventral cranial fissures in osteichthyans. In *Problems in Vertebrate Evolution*, Andrews, S.M., Miles, R.S. & Walker, A.D., eds, Academic Press, London, pp. 227-245.
- GIFFIN, E., 1980. Devonian vertebrates from Australia. *Postilla* 80: 1-15.
- GROSS, W., 1973. Kleinschuppen, Flossenstacheln und Zähne von Fischen aus europäischen und nordamerikanischen Bonebeds des Devons. *Palaeontographica* A 142: 51-155.
- HILLS, E.S., 1936. On certain endocranial structures in *Coccoosteus*. *Geological Magazine* 73: 213-226.
- LINDLEY, I.D. 2002. Lower Devonian ischnacanthid fish (Gnathostomata: Acanthodii) from the Taemas Limestone, Lake Burrinjuck, New South Wales. *Alcheringa* 25(3): 269-291.
- LONG, J.A., 1984a. New placoderm fishes from the Early Devonian Buchan Group, eastern Victoria. *Proceedings of the Royal Society of Victoria* 96(4): 173-186.
- LONG, J.A., 1984b. The plethora of placoderms: the first vertebrates with jaws? In *Vertebrate Zoogeography and Evolution in Australasia*, Archer, M. & Clayton, G., eds, Hesperian Press, Perth, pp. 235-255.
- LONG, J.A., 1988. New palaeoniscoid fishes from the Late Devonian and Early Carboniferous of Victoria. *Memoirs of the Association of Australasian Palaeontologists* 7: 1-64.
- LONG, J.A., 1991. Arthrodire predation by *Onychodus* (Pisces, Crossopterygii) from the Upper Devonian Gogo Formation, Western Australia. *Records of the Western Australian Museum* 15: 369-371.
- LONG, J.A., 1995. *The Rise of Fishes*. UNSW Press, Sydney, 223 pp.
- LONG, J.A. & YOUNG, G.C., 1988. Acanthothoracid remains from the Early Devonian of New South Wales, including a complete sclerotic capsule and pelvic girdle. *Memoirs of the Association of Australasian Palaeontologists* 7: 65-80.
- MAWSON, R., 1987. Early Devonian conodont faunas

- from Buchan and Bindi, Victoria, Australia. *Palaeontology* 30: 251-297.
- MAWSON, R. & TALENT, J.A., 1994. Age of an Early Devonian carbonate fan and isolated limestone clasts and megaclasts, east-central Victoria. *Proceedings of the Royal Society of Victoria* 106: 31-70.
- MAWSON, R., TALENT, J.A., BEAR, V.C., BENSON, D.S., BROCK, G.A., FARRELL, J.A., HYLAND, K.A., PYEMONT, B.D., SLOAN, T.R., SORENTINO, L., STEWART, M.I., TROTTER, J.A., WILSON, G.A. & SIMPSON, A.G., 1988. Conodont data in relation to resolution of stage and zonal boundaries for the Devonian of Australia. In *Devonian of the World*, McMillan, N.J., Embry, A.F. & Glass, D.J., eds, Canadian Society of Petroleum Geologists Memoir 14, v.III: 485-527.
- MAWSON, R., TALENT, J.A., BROCK, G.A. & ENGELBRETSSEN, M.J., 1992. Conodont data in relation to sequences about the Pragian-Emsian boundary (Early Devonian) in south-eastern Australia. *Proceedings of the Royal Society of Victoria* 104: 23-56.
- MCCOY, F., 1876. *Asterolepis ornata* (Eichwald) var. *australis* (McCoy). *Geological Survey of Victoria, Prodromus of the Palaeontology of Victoria* 4: 19-20.
- ØRVIG, T. 1957. Remarks on the vertebrate fauna of the Lower Upper Devonian of Eseuminac Bay, P.Q., Canada, with special reference to the Porolepiform Crossopterygians. *Arkiv för Zoologi* 10(6): 367-426.
- ØRVIG, T., 1969. Vertebrates from the Wood Bay Group and the position of the Emsian-Eifelian boundary in the Devonian of Vestspitsbergen. *Lethaia* 2: 273-328.
- PHILIP, G.M., 1965. Lower Devonian conodonts from the Tyers area, Gippsland, Victoria. *Proceedings of the Royal Society of Victoria* 75: 95-117.
- PICKETT, J.W., TURNER, S. & MYERS, B., 1985. The age of marine sediments near Tumblong, southwest of Gundagai. *Geological Survey of NSW, Quarterly Notes* 58: 12-15.
- POLTIG, W., 1984. Fischreste aus dem Unterdevon von Graz (Steiermark). *Mitteilungen naturwissenschaftlicher vereinfür Steiermark* 114: 107-131.
- SCHULTZE, H.P., 1968. Palaeoniscoidea-Schuppen aus dem Unterdevon Australiens und Kanadas und aus dem Mitteldevon Spitsbergens. *Bulletin of the British Museum of Natural History (Geology)* 16: 343-368.
- SHERWIN, L., 1996. Narramin 1:250000 Geological Sheet SI/55-3: Explanatory notes. Geological Survey of New South Wales, 104pp, Sydney.
- STENSIÖ, E.A. 1945. On the heads of certain Arthrodires II. *Kunglig svenska Vetenskapsakademiens Handlingar* 22: 1-70.
- TALENT, J.A., 1995. Pre-conference excursion: Ordovician-Devonian of southeastern Australia, excursion guide. Auscos-1/Boucot Symposium, pre-conference excursion. 8-17 July 1995. Macquarie University Centre for Ecostratigraphy and Palaeobiology.
- TEICHERT, C. & TALENT, J.A., 1958. Geology of the Buchan area, East Gippsland. *Memoir of the Geological Survey of Victoria* 21: 1-56.
- THOMSON, K.S. & CAMPBELL, K.S.W., 1971. The structure and relationships of the primitive Devonian lungfish *Dipnorhynchus sussmilchii* Etheridge. *Bulletin of the Peabody Museum of Natural History* 38: 1-109.
- TRINAJSTIC, K.M., 1999a. Scales of palaeoniscoid fishes (Osteichthyes: Actinopterygii) from the Late Devonian of Western Australia. *Records of the Western Australian Museum Supplement* 57: 93-106.
- TRINAJSTIC, K. 1999b. Scale morphology of the Late Devonian palaeoniscoid *Moythomasia durgaringa* Gardiner and Bartram, 1977. *Alcheringa* 23: 9-19.
- TURNER, S., 1982. Middle Palaeozoic clasmobranch remains from Australia. *Journal of Vertebrate Paleontology* 2: 117-131.
- TURNER, S., 1991. Palaeozoic vertebrate microfossils in Australasia. In *Vertebrate Palaeontology of Australasia*, Vickers-Rich, P., Monaghan, J.M., Baird, R.F. & Rich, T.H., eds, Pioneer Design Studio & Monash University, Clayton, pp. 429-464.
- TURNER, S., 1993. Palaeozoic microvertebrates of Eastern Gondwana. In *Palaeozoic vertebrate biostratigraphy and biogeography*, Long, J.A., ed, Belhaven Press, London, pp 174-207.
- TURNER, S. & MURPHY, M.A., 1988. Early Devonian microfossils from the Simpson Park Range, Eureka County, Nevada. *Journal of Paleontology* 62: 959-964.
- TURNER, S. & BURROW, C.J., 2000. Annotations to the Devonian correlation Table, B705di00-



- B705ds00: Microvertebrate zonations of East Gondwana. In Weddige, K., ed., *Devon-Korrelationstabelle. Senckenbergiana lethaea* 80(2): 761-763, Frankfurt am Main.
- TURNER, S., WANG S.-T. & YOUNG, G.C., 1995. Lower Devonian microvertebrates from Longmenshan, Sichuan, China: preliminary report. In Lelièvre, H., Wenz, S., Blicek, A. & Cloutier, R. (eds), *Premiers vertébrés et vertébrés inférieurs. Geobios M. S.* 19: 383-387.
- TURNER, S., BASDEN, A.M. & BURROW, C.J., 2000. Devonian vertebrates of Queensland. In *Palaeozoic Vertebrate Biochronology and Global Marine/Non-marine Correlation - Final Report of IGCP 328 (1991-1996)*, Blicek, A. & Turner, S., eds, *Courier Forschungsinstitut Senckenberg* 223: 487-521.
- VALIUKEVICIUS, Y.Y., 1979. Acanthodian scales from the Eifelian of Spitsbergen. *Paleontological Journal* 13: 482-492.
- VALIUKEVICIUS, Y.Y., 1985. Acanthodians from the Narva Regional Stage of the Main Devonian Field. Mosklas, 164pp, Vilnius. (Russian with English summary).
- VALIUKEVICIUS, Y.Y., 1994. Acanthodians and their stratigraphical significance. In *Stratigrafiya i Fauna niznedevonskikh otlojenniy Taryeiskogo opornogo razreza Tcherkesova*, S.V., Karatajute-Talimaa, V.N. & Matukhin, R.G., eds, (Taimyr), pp. 131-197.
- VALIUKEVICIUS, Y.Y., 1998. Acanthodians and zonal stratigraphy of Lower and Middle Devonian in East Baltic and Byelorussia. *Palaeontographica A* 248: 1-53.
- VIETH, J., 1980. Thelodontier-, Acanthodier- und Elasmobranchier-Schuppen aus dem Unter-Devon der Kanadischen Arktis (Agnatha, Pisces). *Göttinger Arbeiten Geologie und Paläontologie* 23: 1-69.
- VIETH-SCHREINER, J. (1983). Fisch-Schuppen und -Zähne aus der Eifeler Kalmulden-Zone (Emsium, Eifelium). *Senckenbergiana lethaea* 64: 129-177.
- WANG N.-Z., 1984. Thelodont, acanthodian and chondrichthyan fossils from the Lower Devonian of southwest China. *Proceedings of the Linnean Society of New South Wales* 107: 419-441.
- WANG N.-Z., 1992. Microremains of agnathans and fishes from Lower Devonian of Central Guangxi with correlation of Lower Devonian between Central Guangxi and Eastern Yunnan, South China. *Acta Palaontologica Sinica* 31(3): 298-307.
- WEBBY, B.D., STEARN, C.W. & ZHEN, Y.Y., 1993. Lower Devonian (Pragian-Emsian) stromatopores from Victoria. *Proceedings of the Royal Society of Victoria* 105: 113-186.
- WELLS, J.W., 1944. Fish remains from the Middle Devonian Bone Beds of the Cincinnati Arch Region. *Palaeontographica Americana* 111: 103-160.
- WHITE, E.I., 1952. Australian arthrodires. *Bulletin of the British Museum of Natural History (Geology)* 1: 249-304.
- WHITE, E.I., 1978. The larger arthrodiran fishes from the area of the Burrinjuck Dam, NSW. *Transcriptions of the Zoological Society of London* 34: 149-262.
- WHITE, E.I. & TOOMBS, H.A., 1972. The buchanoideid arthrodires of Australia. *Bulletin of the British Museum of Natural History (Geology)* 22: 379-419.
- WINCHESTER-SEETO, T., 1996. Emsian Chitinozoa from the Buchan area of southeastern Australia. *Acta Palaontologica Polonica* 41(2): 149-230.
- YOUNG, G.C., 1979. New information on the structure and relationships of *Buchanosteus* (Placodermi: Euarthrodira) from the Early Devonian of New South Wales. *Zoological Journal of the Linnean Society* 66: 309-352.
- YOUNG, G.C., 1993. Middle Palaeozoic macrovertebrate biostratigraphy of eastern Gondwana. In *Palaeozoic Vertebrate Biostratigraphy and Biogeography* Long, J.A., ed., Belhaven Press, London, pp 208-251.
- YOUNG, G.C., LELIÈVRE, H. & GOUJET, D., 2001. Primitive jaw structure in an articulated brachythoracid arthrodire (placoderm fish, Early Devonian) from southeastern Australia. *Journal of Vertebrate Paleontology* 21(4): 670-678.
- YOUNG, V.T., 1995. Micro-remains from Early and Middle Devonian acanthodian fishes from the UK and their biostratigraphic possibilities. *Ichthyolith Issues Special Publication* 1: 65-68; Socorro, New Mexico.
- ZHU M., 2000. Catalogue of Devonian vertebrates in

China, with notes on bio-events. In *Palaeozoic Vertebrate Biochronology and Global Marine/Non-Marine Correlation - Final Report of IGCP 328 (1991-1996)*, Blicek, A. & Turner, S., eds, Courier Forschungsinstitut Senckenberg 223: 373-390.

ZHU M., WANG N.-Z. & WANG J.-Q., 2000. Devonian

macro- and microvertebrate assemblages of China. In *Palaeozoic Vertebrate Biochronology and Global Marine/Non-Marine Correlation - Final Report of IGCP 328 (1991-1996)*, Blicek, A. & Turner, S., eds, Courier Forschungsinstitut Senckenberg 223: 361-372.

Manuscript received 13 June 2002

Revision accepted 5 April 2003

TWO ALPINE WOLF SPIDERS OF AUSTRALIA: *ARTORIA ALTA* SP.  
NOV., AND THE MALE OF *LYCOSA MUSGRAVEI* MCKAY, 1974  
(ARANEAE, LYCOSIDAE)

VOLKER W. FRAMENAU

Department of Zoology, University of Melbourne, Parkville, Victoria 3010  
Current address: Department of Terrestrial Invertebrates, Western Australian Museum, Francis St, Perth,  
Western Australia 6000

FRAMENAU, V. W., 2003; 06:15. Two Alpine Wolf Spiders of Australia: *Artoria alta* sp. nov., and the male of *Lycosa musgravei* McKay, 1974 (Araneae, Lycosidae). *Proceedings of the Royal Society of Victoria* 115(1): 27-34. ISSN 0035-9211

A new wolf spider species from the alpine region of the Australian Alps, *Artoria alta* sp. nov., is described and the male of *Lycosa musgravei* McKay, 1974 is illustrated for the first time. Dense pubescence in *L. musgravei* may have evolved as an adaptation to the alpine environment to insulate against heat loss and protect against high levels of radiation.

*Key words:* Wolf Spiders, *Artoria alta*, *Lycosa musgravei*, Australian Alps, pubescence

THE ALTITUDINAL PROFILE of the Australian Alps can be divided into three main ecological zones: alpine, subalpine, and montane (McLuckie & Petric 1927). The alpine zone includes the area above the physiological limit of tree growth (ca. 1800 m). In the Australian Alps, it has a continuous snow cover for at least four months per year, and for six to eight months the minimum temperature is below freezing. Precipitation is high and ranges from 1,800 – 3,100 mm per year, with about 60% falling as snow (Costin 1957). Herbfields and heathlands dominate the vegetation and approximately 200 species in at least ten different plant communities are recognized (Costin *et al.* 1979). Globally, the alpine zone has its upper limit where there is a permanent cover of snow or ice (nival zone); however, the temperature regime even at its highest elevation at Mt Kosciuszko (2,230m) is too warm to establish a nival zone in the Australian Alps. The subalpine zone reaches from the treeline down to the winter snowline, signified by the presence of a continuous cover of snow for at least one month (ca. 1500 m). Minimum mean temperatures below freezing prevail for about six months, and precipitation ranges from ca. 770 – 2200 mm. Snow Gum (*Eucalyptus pauciflora*) dominates the woodland, in addition to wet and dry heathland and sod tussock grassland. The subalpine zone changes into the montane zone (ca. 900 – 1500 m), with the transition from Snow Gum woodland to eucalypt forest (Costin 1975).

Despite a short vegetation period and comparatively harsh conditions, the alpine and subalpine environment host a diverse fauna. In addition to species which extend their range from lower altitudes only in the favourable summer months, some animals facultatively live above the treeline ('eualpine') or at least may be able to complete their whole life cycle there ('tyehoalpine') (Hesse 1924; 'alpine residents' in Green & Osborne 1994). Invertebrates, in particular grasshoppers and spiders, show a high number of alpine species in the Australian Alps (Green & Osborne 1994). The diversity of spiders appears to be unaffected by the change from a woodland to treeless alpine environment. About 20 families have been reported to occur in alpine and subalpine regions of the Australian Alps but a large number of species remain undescribed (Green 1988; in Green & Osborne 1994). Wolf spiders (Lycosidae) belong to the better-known taxa, and some appear to be particularly abundant in tall alpine herbfields and tussock grasslands. Three species of lycosids from the alpine region of Mt Kosciuszko have been scientifically named: *Lycosa kosciuskoensis* McKay, 1974, *L. summa* McKay, 1974, and *L. musgravei* McKay, 1974, all known from females only. Other lycosids may be found at high altitudes but appear to occur mainly in forests of the montane region, such as *Venatrix finesta* (C. L. Koeh, 1847) and *V. australiensis* Framenau & Vink, 2001 (Framenau & Vink 2001).



A number of morphological adaptations found in high altitude arthropods are attributed to alpine environmental conditions (Mani 1968; Sømme 1989): Increased hairiness (pubescence) is thought to serve as thermal insulation and protection against ultra-violet light; increased melanism (darkening) may improve heat absorption; and a decrease in size is thought to be due to a shorter time for growth, reduced food availability, and increased shelter options. For example, the Central European wolf spider *Pardosa saturator* Simon, 1937, which is found above ca. 800 m is darker in colouration than its sibling species *P. wagleri* (Hahn, 1822), which mainly occurs at lower altitudes (Barthel & von Helversen 1990; Manderbach & Framenau 2001). However, *P. saturator* is significantly larger than *P. wagleri*, contradicting the predictions of Mani (1968) and Sømme (1989).

Recent examinations of the collections of the Australian Museum, Sydney, the Australian National Insect Collection, Canberra, and the Museum Victoria, Melbourne, provided new wolf spider material from the alpine zone of the Australian Alps. The aim of this study is to facilitate the identification and study of alpine wolf spiders by describing a new species, *Artoria alta* sp. nov., and illustrating the male of *L. musgravei* of which only the holotype female was known (McKay 1974). In addition, some morphological features of these species are discussed as adaptations to the extreme conditions of the alpine environment.

## MATERIALS AND METHODS

Descriptions are based on specimens preserved in 70% ethanol. The epigyne of a female *L. musgravei* was prepared for examination by submersion in 10% KOH overnight at room temperature. For clarity, the illustrations of male and female genitalic organs omit the setae. The morphological nomenclature follows Dondale & Redner (1990), Framenau & Vink (2001), and Framenau (2002).

### Abbreviations

Eyes. Anterior (AE), anterior median (AME), anterior lateral (ALE), posterior (PE), posterior median (PME), posterior lateral (PLE). Measurements: total length (TL), carapace length (CL) and width (CW), abdomen length (AL) and width (AW). Genitalia: Male pedipalp: embolus (E), basoembolic apophysis (BEA), median apophysis (MA), palpa (PA), subtegulum (STE), tegulum (TEG), terminal apophysis (TA). Female genitalia: copulatory duet (CD),

median septum (MS), posterior transverse part (PTP), spermatheca (SP).

### Collections

Australian Museum, Sydney (AM); Australian National Insect Collection, Canberra (ANIC); Museum of Victoria, Melbourne (MV).

## SYSTEMATICS

### *Artoria alta*, sp. nov.

Figs. 1A-D, 2

*Material examined.* Holotype. ♀, New South Wales, Mt Koseiuko NP, near Smiggin Holes, 1700 m, alpine moor, 36°24'S, 148°26'E, 7.xii.1994, coll. Daniel Bickel (AM KS44789).

*Paratypes.* 1 ♀, 1 immature ♀, New South Wales, Mt Koseiuko NP, Spence Cr near Charlottes Pass, 36°24'S, 148°21'E, 28.xi.1994, coll. Daniel Bickel (AM KS45825).

*Diagnosis.* The shape of the MA of the male pedipalp of *A. alta* that has a base with two distinct grooves and a triangular apical part is unique within the genus *Artoria*.

### Description

#### Males

*Carapace* (Fig. 1A). Brown, with a distinct light brown median band narrowing posteriorly, constricted anteriorly of fovea, and with a narrow black median line in head region; indistinct light brown submarginal bands; dark grey radial pattern; carapace covered with short black setae except in median band; two rows of black bristles between PME, one long black bristle between AME, and two long bristles below AE. *Sternum*: Uniformly dark brown and sparsely covered with brown setae and bristles, less dense centrally. *Labium*: Brown, basally dark brown; front end truncate and white. *Chelicerae*: Uniformly dark brown; covered with few white setae and black bristles; three (holotype male four on right) retromarginal teeth, with the median largest, two promarginal teeth, with the apical larger. *Pedipalp* (Figs 1B-D): Cymbium with scopulous setae dorsally in apical half; base of MA with two distinct grooves and its apical part triangular; E long and stout, resting in a groove of the sickle-shaped TA (Fig. 1D). *Abdomen*: Brown; indistinct dark median band with darker lateral borders; lanceolate heart mark in anterior half very distinct as a result of a dense cover of white




Fig. 1. Male holotype (AM KS44789) of *Artoria alta*, sp. nov.; A, male, *habitus*; B, C, left male pedipalp, ventral and retrolateral view; D, left male pedipalp, apical part of bulb (MV KS45825). Scale bar: A, 1.5 mm; B, C, 0.34 mm; D, 0.25 mm. BEA, basocymbolic apophysis; E, embolus; MA, median apophysis; PA, palea; STE, subtegulum; TA, terminal apophysis; TEG, tegulum.

setae; sparsely covered in white setae and few short, black bristles; venter light brown; setae and bristles as dorsally; spinnerets light brown. *Legs*: Leg formula IV > III > I > II; uniformly brown; spination of leg I: Femur: 3 dorsal, 1 apicoprolateral; tibia: 3 ventral pairs; metatarsus: 3 ventral pairs; 2 prolateral.

*Measurements* (based on holotype). TL 5.8, CL 3.0, CW 2.2. Eyes: AME 0.09, ALE 0.08, PME 0.3, PLE 0.22. Row of eyes: AE 0.54, PME 0.80, PLE 1.02. Sternum (length/width) 1.4/1.2. Labium (length/width) 0.20/0.20. AL 2.9, AW 1.9. *Legs*: Lengths of segments (femur + patella/tibia + metatarsus + tarsus = total length): Pedipalp 1.0+0.95+ - +0.8 = 2.75, I 2.9+2.4+1.5+0.95 = 6.75, II 1.85+2.25+1.5+1.0 = 6.6, III 1.8+2.0+1.65+0.85 = 6.3, IV 2.3+3.0+2.65+1.15 = 9.1. *Size variation* (male paratype): TL 5.6; CL 2.7; CW 1.85.

#### Female

Mature female unknown. The colouration of the im-

mature paratype female is similar to the colouration of the male. Its size (TL 6.1, CL 2.55, CW 1.55) suggests a size dimorphism in this species. The female




Fig. 2. Records of *Artoria alta*, sp. nov.



appears to be larger, as reported in all other species of *Artoria* (Framenau 2002).

**Remarks.** The genus *Artoria* Thorell, 1877 was recently revised in part to include 12 Australian species, however, more than 50 species may exist (Framenau 2002). The genus is mainly defined by the presence of a broad, basoembolic apophysis (BEA), and the position (apical on tegulum) and shape (spoon-shaped or strongly bifurcate) of the median apophysis of the male pedipalp. The triangular shape of the apical part of the median apophysis of *A. alta* represents a unique modification of the spoon-shaped form.

The subfamilial division of wolf spiders remains controversial (e.g., Dondale 1986, Zyuzin 1993, Sierwald 2000, Vink *et al.* 2002). Recent molecular analysis supports the uniqueness of Australasian lycosid genera as suggested by distinct morphological features (Vink *et al.* 2002). In particular the male genital morphology of *Artoria*, as well as of the related New Zealand genera *Anoteropsis* L. Koch, 1878 and *Notocosa* Vink, 2002 (see Vink 2002), does not appear to conform to any of the five main lycosid subfamilies established by Dondale (1986), or the additional Evippinac and Wadicosinae (Zyuzin 1985), Piratinae (Zyuzin 1993), or Tricassinac (Alderweireldt & Joequé 1993). However, I regard the erection of a new subfamily for *Artoria*, *Anoteropsis* and *Notocosa* premature without further examining the full morphological variation within this group, in particular within the large number of undescribed *Artoria*.

*Artoria alta*, as inferred from the three specimens examined, does not show any clear morphological adaptations to its alpine habitat. The species is not larger, darker in its colouration (melanism) or has a denser pubescence than any other Australian *Artoria* described (Framenau 2002).

**Distribution.** Only known from Mt Kosciusko, New South Wales (Fig. 2).

**Etymology.** The species name is an adjective in apposition derived from the Latin, *altus*, meaning high, and refers to the high altitude at which this species is found.

*Lycosa musgravei* McKay, 1974  
Figs. 3A-G, 4

*Lycosa musgravei* McKay, 1974: 34-35, Figs 2A-C.-Brignoli, 1983: 450; McKay, 1985: 80.

**Material examined.** Holotype. ?, New South Wales,

Mt Kosciusko, 36°27'S, 148°16'E, 30.i.1966, L. Voysey (AM KS23).

**Other material examined.** **Australian Capital Territory:** 1 female with spiderlings, Mt Gingera, 35°34'S, 148°47'E, 28.v.1970, coll. M. S. Upton, 5800 ft, from silk lined vertical burrow (ANIC); 1 female, Mt Gingera, 35°34'S, 148°47'E, 28.v.1970, coll. M. S. Upton, from silk lined vertical burrow (ANIC); 1 male, Mt Gingera, 35°34'S, 148°47'E, 28.iii.1970, coll. M. S. Upton, 5800ft, free ranging on snow (ANIC). **New South Wales:** 1 female, Bombala, 36°54'S, 149°14'E, i.1930, coll. A. J. Barrett (AM KS84075); 1 male, Charlotte Pass, Mt Kosciusko, 36°24'S, 148°19'E, 8.v.1980, coll. J. Balderson (ANIC); 1 male, 1 female, Crackenback Peak, 36°24'S, 148°32'E, iii.1964, coll. W. A. Howard (ANIC); 2 females, Gungahlin River (tributary of Snowy River), 36°17'S, 148°52'E, no date, coll. R. J. Jenner (AM KS84081); 1 female, Kiandra, 35°52'S, 148°29'E, 21.ii.1960, coll. E. F. Rick (ANIC); 1 female, Mt Kosciusko, 36°27'S, 148°15'E, no date, 6000ft (AM KS84082); 1 male, Mt Kosciusko NP, Wraggs Ck/Pipers Creek Aqueduct, 36°32'S, 148°28'E, 31.iii.1982 (ANIC); 1 female, Perisher Range, East of Guthega Dam, 36°25'S, 148°25'E, 28.i.1962, coll. K. Horne, 5500ft, in burrow (AM KS84079); 1 female, Spenceers Creek, Mt Kosciusko, 36°27'S, 148°16'E, 22.xi.1952, coll. A. Musgrave, 1000m (AM KS70010); 1 female, Spenceers Creek, Mt Kosciusko, 36°27'S, 148°16'E, 24.xi.1952, coll. A. Musgrave, 1000m (AM KS70012); 1 female, Spenceers Creek, Mt Kosciusko, 36°27'S, 148°16'E, 23.xi.1952, coll. C. E. Chadwick, 1000m (AM KS70013); 1 female, Tumut, 35°18'S, 148°13'E, 7.iii.1949, coll. S. Bayliss (AM KS84080); 1 female, Tumut Pond, 35°18'S, 148°13'E, 20.xii.1951, coll. K. R. Sharp (AM KS82593). **Victoria:** 2 females, Elsternwick, 37°53'S, 145°00'E, 20.i.1954, coll. Mr Brownlie (MV K8095, K8209); 3 females, Hotham Heights, 36°59'S, 147°08'E, xii.1933, coll. A. Musgrave (AM KS84076, KS84078); 1 male, Mt Bogong, 36°44'S, 147°18'E, 21.ii.1972, coll. C. Kohlman (MV K8204); 1 male, Mt Bogong, 36°44'S, 147°18'E, 21.ii.1972, coll. C. Kohlman (MV K8207); 1 female, Mt Buffalo, 36°46'S, 146°46'E, xii.1933, coll. A. Musgrave, 4000ft (AM KS84077); 1 female with eggsae (106 larvae, 71 undeveloped eggs), Mt Gibbo, 36°36'S, 147°57'E, 15.i.1975 (MV K8094); 1 female, Mt Hotham, 36°59'S, 147°08'E, i.1945, 6000ft (MV K8205); 1 female, 1 male, Mt Hotham, summit,




Fig. 3. Male (MV K8204) and female (MV K8095) of *Lycosa musgravei* McKay. A, male, *habitus*; B, carapace, lateral view; C, D, left male pedipalp, ventral and retrolateral view; E, left male pedipalp, apical section of bulbus; F, G, female epigyne, ventral and dorsal view. Scale bar: A, 4.6 mm; B, 6.2 mm; C, D 1.58 mm; E, 1.24 mm; F, G, 1.67 mm. CD, copulatory ducts; E, embolus; MA, median apophysis; MS, median septum of epigyne; PA, palca; PTP, posterior transverse part; SP, spermatheca; STE, subtegulum; TA, terminal apophysis; TEG, tegulum.

36°59'S, 147°08'E, 27.i.1973, coll. M. Pearce, 6100ft, (MV K8206, K8208); 1 female, North Mt Phipps, 1.5km SW Moe, Head of Spring Ck, 37°12'S, 147°26'E, 28.xii.1989, coll. Heath Morris, 4000ft (MV K8166).

**Diagnosis.** *Lycosa musgravei* is similar in general appearance to *L. gilberta* Hogg, 1905, *L. godeffroyi* L. Koch, 1865 and *L. leuckartii* (Thorell, 1870), three species commonly found in lowland areas of southern and south-eastern Australia (Queensland, New South Wales, Victoria, South Australia, and Western Australia). Males can be distinguished by the shape of the TA, which is double-lamellar in *L. musgravei*, but simply sickle-shaped in *L. godeffroyi* and which forms a wave-like structure in *L. gilberta* and *L. leuckartii*. The lateral tips of the PTP of the female epigyne are strongly curved anteriorly in *L. musgravei*, but point laterally in *L. gilberta*, *L. godeffroyi* and *L. leuckartii*. The abdominal pattern of *L. musgravei* differs dis-

tinctly from that of the other two large, alpine lycosids, *L. kosciuskoensis* ('dark brown longitudinal spot surrounded by narrow fawn band that becomes somewhat




Fig. 4. Records of *Lycosa musgravei* McKay.



diffuse posteriorly') and *L. summa* ('pale brown to fawn (...) longitudinal stripe (...) encloses a dark brown hastate stripe anteriorly') (descriptions cited after McKay 1974; also pers. observation).

### Description

#### Males

**Carapace** (Fig. 3A-B). Dorsal line in lateral view straight from head region to fovea, then abruptly descending (Fig. 3B); dark reddish brown with light brown median band and four white, but dark edged, radial stripes, the anterior ones reaching below PLE; indistinct light brown submarginal bands; carapace densely covered with short black setae with the exception of head region, median band and light radial stripes which are densely covered with white setae; black bristles in head region and lateral of PLE; one long, black bristle between AME and four long bristles below AE. **Sternum**: Light brown, densely covered with black setae; fewer bristles increasing in length towards margins. **Labium**: Dark brown, front end truncate and light. **Chelicerae**: Reddish brown; densely covered with white setae and fewer black bristles; three retromarginal teeth of similar size, three promarginal teeth with the median largest and the two apical ones fused at the base. **Pedipalp** (Figs 1C-E): Lateral tip of MA with little apical teeth; E broad over all of its length; TA a double lamellar structure with its tip bent basally (Fig. 1E). **Abdomen**: Brown; light brown median band in anterior half contains a double wedge-shaped, dark marking with elongated, black posterior ends; four to five dark chevrons in posterior half; densely covered with setae of variable colour corresponding to abdomen colouration; lateral light brown with dense cover of white setae; venter black with dense cover of black setae; spinnerets dark brown. **Legs**: Leg formula IV > III > I > II; uniformly brown; dense, scopulous setae on all tarsi, metatarsi I and II, and in apical half of metatarsus III and apical third of metatarsus IV; spination of leg I: Femur: 3 dorsal, 2 apicoprolateral, 2 retrolateral; patella: 1 prolateral, 1 retrolateral; tibia: 2 dorsal; 3 ventral pairs, 2 prolateral, 2 retrolateral; metatarsus: 2 ventral pairs; 2 prolateral, 2 retrolateral, 2 small apicoprolateral; 2 small apicoretrolateral.

**Measurements** (based on MV K8204). TL 18.0, CL 9.9, CW 7.3. Eyes: AME 0.34, ALE 0.35, PME 0.7, PLE 0.60. Row of eyes: AE 1.8, PME 1.9, PLE 2.4.

Sternum (length/width) 4.2/3.0. Labium (length/width) 1.1/1.3. AL 3.0, AW 1.8. Legs: Lengths of segments (femur + patella/tibia + metatarsus + tarsus = total length): Pedipalp 3.5+3.4+ +3.8 = 10.7, I 8.0+10.2+7.2+4.1 = 29.5, II 7.4+8.8+6.7+4.0 = 26.9, III 6.5+7.5+6.3+3.7 = 24.0, IV 8.1+10.3+8.7+4.5 = 31.6. **Size variation** (range, mean  $\pm$  SE): TL 19.0 – 20.6, 19.6  $\pm$  0.9; CL 9.9 – 11.5, 10.7  $\pm$  0.8; CW 7.3 – 8.3, 8.1  $\pm$  0.7; n = 3.

#### Females

The female of *L. musgravei* is described in detail in McKay (1974). Cephalothorax colouration agrees with that of the male, the abdominal pattern is less distinct. The epigyne of a specimen collected in Victoria is depicted here in ventral (Fig. 3E) and dorsal view (Fig. 3F) to illustrate diagnostic features. Size variation of the material deposited in the MV is given, as only the holotype female was previously known (McKay 1974). **Size variation** (range, mean  $\pm$  SE): TL 21.0 – 28.5, 24.3  $\pm$  2.8, n = 5; CL 11.0 – 15.0, 12.9  $\pm$  1.3, n = 6; CW 8.3 – 11.1, 9.6  $\pm$  1.1; n = 6.

**Remarks** *Lycosa* Latreille, 1804 has recently been suggested to be an exclusively Mediterranean genus (Zyuzin & Logunov 2000). In addition, preliminary molecular data suggests that some of the large burrowing lyeosids of Australia, of which *L. musgravei* is part of and which were represented in their analysis by *Lycosa godeffroyi*, is close to the North American genus *Geolycosa* Montgomery (Vink et al. 2002). However, the generic description of *Geolycosa* does not match the Australian species (e.g. Dondale & Redner 1990): the dorsal profile of the carapace of *Geolycosa* is highest in the cephalothorax region, followed by a gentle slope towards the posterior cephalothorax margin, whereas the dorsal profile of *L. musgravei* is straight over most of the carapace and descends sharply behind the fovea (Fig. 3B). In addition, *Geolycosa* does not have light median and submarginal bands and a Union-Jack-pattern of white radial bands on the carapace which are present in *L. musgravei*, *L. gilberta*, *L. godeffroyi* and *L. leuckartii*. Three recent revisions of Australasian lyeosid genera, *Allotrochosina* Roewer, 1960, *Artoria* Thorell, and *Venatrix* Roewer, 1960, suggest the uniqueness of the Australasian fauna (Vink 2001; Framenau & Vink 2001; Framenau 2002). Likewise, *L. musgravei*, and all other Australian lyeosids placed in *Lycosa*, almost certainly belong to different, most likely new genera; however, *L. musgravei* is retained in *Lycosa* pending a full generic revision of Australian wolf spiders, which is currently



being conducted by the author at the Western Australian Museum.

The whole body of *L. musgravei* is covered with a very dense layer of setae and bristles, which are much denser than in any of the specimens of the related species *L. gilberta*, *L. godeffroyi* or *L. leuckartii*, (pers. observation). This dense pubescence conforms to one of the adaptations of high altitude arthropods to insulate against heat loss and protect against high levels of radiation (Mani 1968, Somme 1989).

**Habitat and phenology.** *Lycosa musgravei* is reported to be the most frequently observed wolf spider in the Snowy Mountains, with their open burrow found in tall alpine herbfields and grasslands (Green & Osborne 1994). The burrow (closed with a thick sheet of webbing in winter) is raised above ground level by inclusion of an open turret of web and vegetation litter (Green & Osborne 1994). The records of two females from Elsternwick (suburban Melbourne) seem to be unusual for this apparently alpine spider, and may be the result of human recreational activities.

Mature females were found from November through to May with the exception of April, with most records in December and January. One female with eggsae and a female carrying young were collected in December and May respectively. Mature males were found between January and May, with the exception of April as in females.

#### Distribution

Australian Capital Territory, New South Wales and Victoria (Fig. 4).

#### ACKNOWLEDGEMENTS

I appreciated the support of Ken Walker, Catriona McPhee (both MV), Mike Gray, and Graham Milledge (both AM) for their assistance in investigating the lycosid collections under their care. I am also indebted to Bruce Halliday (ANIC) for the loan of material. Ken Green and David Rowell assisted in the (unfortunately unsuccessful) search for male specimens of *L. kosciuskoensis* and *L. summa*. Mark Elgar provided laboratory facilities, David Paul digitalised the drawings, and Melissa Thomas and Mark Harvey provided helpful suggestions to improve earlier drafts of this manuscript.

#### REFERENCES

- ALDERWEIRELDT, M. & JOCQUÉ, R., 1993. A redescription of *Tricassa deserticola* Simon, 1910, representing the Tricassinae, a new subfamily of wolf spiders (Araneae, Lyeosidae). *Belgian Journal of Zoology* 123: 27-38.
- BARTHEL, J. & VON HELVERSEN, O., 1990. *Pardosa wagleri* (Hahn 1822) and *Pardosa saturator* Simon 1937, a pair of sibling species (Araneae, Lyeosidae). *Bulletin de la Société Européenne d'Arachnologie* 1: 17-23.
- BRIGNOLI, P. M., 1983. *A catalogue of the Araneae described between 1940 and 1981*. Manchester University Press in association with The British Arachnological Society, Manchester, 755 pp.
- COSTIN, A. B., 1957. The high mountain vegetation of Australia. *Australian Journal of Botany* 5: 173-189.
- COSTIN, A. B., 1975. Sub-alpine and alpine communities. In *Australian Grasslands*, R. M. Moore, ed., Australian National University Press, Canberra, 191-198.
- COSTIN, A. B., GRAY, M., TOTTERDELL, C. J. & WIMBUSCH, D. J. 1979. Kosciusko Alpine Flora, CSIRO, Melbourne, 408 pp.
- DONDALE, C. D., 1986. The subfamilies of wolf spiders (Araneae: Lyeosidae). *Actas X Congreso Internacional de Aracnología, Jaca, España* 1: 327-332.
- DONDALE, C. D. & REDNER, J. H., 1990. The wolf spiders, nurseryweb spiders, and lynx spiders of Canada and Alaska. Araneae: Lyeosidae, Pisauridae, and Oxyopidae. *The Insects and Arachnids of Canada* 17: 1-383.
- FRAMENAU, V. W. & VINK, C. J., 2001. Revision of the wolf spider genus *Venatrix* Roewer (Araneae, Lyeosidae). *Invertebrate Taxonomy* 15: 927-970.
- FRAMENAU, V. W., 2002. Review of the wolf spider genus *Artoria* Thorell (Araneae: Lyeosidae). *Invertebrate Systematics* 16: 209-235.
- GREEN, K., 1988. *A Study of Antechinus swainsonii and Antechinus stuartii and their Prey in the Snowy Mountains*. Ph.D. thesis, Zoology Department, Australian National University.
- GREEN, K. & OSBORNE, W., 1994. *Wildlife of the Australian Snow-Country*. Reed Books, Chatswood (NSW), 200 pp.
- HAHN, C. W., 1822. *Monographie der Spinnen. Heft*

3. Nürnberg: 1-2, 4 plates.
- HOGG, H. R., 1905. On some South Australian spiders of the family Lycosidae. *Proceedings of the Zoological Society London* 2: 569-590.
- HESSE, R., 1924. *Tiergeographie auf ökologischer Grundlage*. Gustav Fischer Verlag, Jena, 613 pp.
- KOCH, C. L., 1847. *Die Arachniden. 14. Band*. Zehsche Buchhandlung, Nürnberg: 89-210.
- KOCH, L., 1865. Beschreibungen neuer Arachniden und Myriapoden. *Verhandlungen der zoologisch-botanischen Gesellschaft Wien* 15: 857-892.
- KOCH, L., 1878. *Die Arachniden Australiens*. Bauer and Raspe, Nürnberg: 969-1044.
- Latreille, P. A., 1804. Tableau méthodique des insectes. *Nonveau Dictionnaire d'Histoire Naturelle*, Paris 24: 129-295.
- MANDERBACH, R. & FRAMENAU, V. W., 2001. Spider (Arachnida: Araneae) communities of riparian gravel banks in the northern parts of the European Alps. *Bulletin of the British arachnological Society* 12: 1-9.
- MANI, M. S., 1968. *Ecology and Biogeography of High Altitude Insects*. W. Junk, The Hague, 527 pp.
- McKAY, R. J., 1974. The wolf spiders of Australia (Araneae: Lycosidae): 4. Three new species from Mount Kosciuszko, N.S.W. *Memoirs of the Queensland Museum* 17: 27-36.
- McKAY, R. J., 1985. Lycosidae. In *Zoological Catalogue of Australia, Vol. 3. Arachnida, Mygalomorphae, Araneomorphae in Part, Pseudoscorpionida, Amblypygida, Palpigradi*, D. W. Walton, ed., Australian Government Publishing Service, Canberra, 73-88.
- McLUCKIE, J. & PETRIE, H. K., 1927. The vegetation of the Kosciusko Plateau. Part. 1. The plant communities. *Proceedings of the Linnean Society of New South Wales* 52: 187-221.
- ROEWER, C. F., 1960 [imprint date 1959]. Araneae Lycosaeformia II (Lycosidae) (Fortsetzung und Schluss). *Exploration du Parc National de l'Upemba - Mission G. F. de Witte* 55: 519-1040.
- SIERWALD, P., 2000. Description of the male of *Sosippus placidus*, with notes on the subfamily Sosippinae (Araneae, Lycosidae). *Journal of Arachnology* 28: 133-140.
- SIMON, E., 1937. *Les arachnides des France. Tome VI. Synopsis générale et catalogue des espèces françaises des l'ordre des Araneae; 5e et dernière partie*. Paris, 6: 979-1298.
- SOMME, L., 1989. Adaptations of terrestrial arthropods to the alpine environment. *Biological Reviews* 64: 367-407.
- THORELL, T., 1870. Araneae nonnullae Novae Hollandie, descriptae. *Öfversigt af Kongliga Svenska Vetenskaps-Akademien Förhandlingar* 27: 367-389.
- THORELL, T., 1877. Studi sui ragni Malesi e Papuani. I Ragni di Selebes raccolti nel 1874 dal Dott. O. Beccari. *Annali di Museo Civico di Storia Naturale Genova* 10: 341-634.
- VINK, C. J., 2001. A revision of the genus *Allotrochosina* Roewer (Araneae: Lycosidae). *Invertebrate Taxonomy* 15: 461-466.
- VINK, C. J., 2002. Lycosidae (Arachnida: Araneae). *Fauna of New Zealand*, Manaaki Whenua Press, Lincoln, New Zealand.
- VINK, C. J., MITCHELL, A. D. & PATERSON, A. M. 2002. A preliminary molecular analysis of phylogenetic relationships of Australasian wolf spider genera (Araneae, Lycosidae). *Journal of Arachnology* (in press).
- ZYUZIN, A. A., 1985. Generic and subfamilial criteria in the systematics of the spider family Lycosidae (Aranei), with the description of a new genus and two new subfamilies. *Trudy Zoologicheskogo Instituta, Akademia Nauk SSSR* 139: 40-51.
- ZYUZIN, A. A., 1993. Studies on the wolf spiders (Araneae: Lycosidae). I. A new genus and new species from Kazakhstan, with comments on the Lycosinae. *Memoirs of the Queensland Museum* 33: 693-700.
- ZYUZIN, A. A. & LOGUNOV, D. V., 2000. New and little known species of the Lycosidae from Azerbaijan, the Caucasus (Araneae, Lycosidae). *Bulletin of the British arachnological Society* 11: 305-319.

Manuscript received 26 June 2002

Revision accepted 14 August 2003



*GLAUCODON BALLARATENSIS* (MARSUPIALIA, DASYURIDAE), A LATE  
PLIOCENE 'DEVIL' FROM BATESFORD, VICTORIA.

W. R. GERDTZ & N. W. ARCHBOLD

School of Ecology and Environment, Melbourne Campus, Deakin University, Burwood, Victoria, 3125

GERDTZ, W. R. & ARCHBOLD, N. W., 2003:11:30. *Glaucodon ballaratensis* (Marsupialia, Dasyuridae), a late Pliocene 'devil' from Batesford, Victoria. *Proceedings of the Royal Society of Victoria* 115(2): 35-44. ISSN 0035-9211.

The right mandible of a dasyurid from Pliocene sediments at Batesford, near Geelong, Victoria is described as a new specimen of *Glaucodon ballaratensis* Stirton, 1957. The new specimen is morphologically close to the holotype of *Glaucodon ballaratensis*. Several dental characteristics of the new specimen unknown from the holotype of *Glaucodon ballaratensis* are close to those of *Sarcophilus moornaensis* Crabb, 1982. *Glaucodon ballaratensis* also shares features with *Dasyurus maculatus* Kerr, 1792 and *Sarcophilus laniarius harrisii* Boitard, 1842, and hence the Batesford specimen offers additional information on the origins of these dasyurids.

*Key words:* *Glaucodon ballaratensis*, Pliocene, Batesford, Moorabool Viaduct Sands, Dasyuromorphia, Dasyuridae.

THE DASYUROMORPHIAN fossil record represents the dominant marsupial insectivore-carnivore radiation in Australasia during the Quaternary. Discoveries from Late Oligocene and Miocene deposits in Riversleigh in northwestern Queensland have extended this record (Wroe, 1998), and Wroe (2003) provides a modern summary of fossil Dasyuromorphian record. However, the order is poorly represented in the Pliocene, where the species *Glaucodon ballaratensis* Stirton, 1957, *Dasyurus dunnalli* Bartholomai, 1971, *Archerium chinchillaensis* Wroe & Maekness 2000 and *Sarcophilus moornaensis* Crabb, 1982 are the only known dasyuromorphians. The Pliocene record may also include indeterminate species of *Antechinus* from Grange Burn, Victoria (Turnbull & Lundelius, 1970), and *Planigale* from Bluff Downs in Queensland (Archer, 1982).

The monotypic genus *Glaucodon* was described by Stirton (1957) and based on a single right mandible with the M4 present. The holotype was found in the Parish Well at Smeaton, near Ballarat. The specimen was originally presented to the National Museum of Victoria in 1914 by Mr. J. Marshall, and remained undescribed for over forty years. E. D. Gill revisited the type location and discovered the first molar (M1) which fits into the alveolus and contacts perfectly with the broken root of the posterior alveolus (Gill, 1953). The holotype of *Glaucodon ballaratensis* (NMV

P16136) therefore includes the remains collected by both Marshall and Gill. In 1964, W. D. Turnbull recognised two fragments in the Museum Victoria Vertebrate Palaeontology Collections as coming from the well at Smeaton that belonged to the holotype of *G. ballaratensis*. Together the fragments provide much of the outline of the ascending ramus from the left dentary. This material was not available to Stirton (1957) when the holotype was first described.

Stirton did not assign *G. ballaratensis* a geological age with any certainty. Gill's analysis of the sediments surrounding the supplementary material collected in 1953 indicated that the specimen's stratigraphical position in the section at Smeaton can be established relative to the overlying basalt flows (Gill, 1953). Another analysis of the Smeaton locality undertaken in 1993 noted that the fossiliferous layer that contained *G. ballaratensis* lies above a basalt flow which correlates closely with a nearby flow with a radiometric date of 2.1 myBP (Turnbull, Lundelius & Tedford, 1993). Basalt flows above the fossiliferous layer have been interpreted as no older than 1.9 myBP (Aziz-ur-Rhman & McDougall, 1972). This implies that the sediment containing *G. ballaratensis* represents an interval between basalt flows of the ages of 1.9 myBP and 2.1 myBP, which implies a Late Pliocene age for the holotype.

The right mandible of a fossil dasyurid with near complete dentition (NMV P207018) was presented to



the Museum of Victoria in 1996 by Mr. P. Robertson. Robertson's specimen was found in the Moorabool Viaduct Sands (Pliocene) outcropping at Portland Cement Limited's Limestone quarry at Batesford, Victoria. The specimen was referred to *G. ballaratensis* (T. H. Rich, pers. comm. 1997, see Gerdzt, 2001). Morphological characteristics of the teeth of this specimen fully confirm that the specimen belongs to this species. Measurements and comparisons of the teeth and mandible of Robertson's specimen reveal differences in relative proportions of the molar trigonids and mandibular depth to those of the holotype, however the differences are considered to be of infraspecific importance.

The significance of *G. ballaratensis* in dasyuromorphian evolution has been discussed by Gill (1953), Stirton (1957), Ride (1964), Marshall (1973), Archer (1976, 1982), Archer and Bartholomai (1978) and Crabb (1982). Characteristics of the Batesford specimen offer additional information on the phylogeny of the Dasyuridae, specifically the relationship of *Glaucodon* to *Dasyurus maculatus* Kerr, 1792, *Sarcophilus laniarius harrisi* Boitard, 1842 and *Sarcophilus moornaensis* Crabb, 1982.

#### STRATIGRAPHY AND AGE OF THE BATESFORD SPECIMEN

It has been suggested (T. H. Rich, pers. comm.) that the distinctive preservation of the Batesford specimen of *G. ballaratensis* indicates that it was originally located in a fissure filling within the argillaceous Moorabool Viaduct Sands of Pliocene age. The unit, as exposed at Batesford Quarry, consists of a black, silty clay with a notable absence of finely comminuted marine fossils (Rich, 1976). The dark brown coloration of the mandible and black coloration of the tooth enamel of the specimen corresponds with other vertebrate remains attributed to this unit, notably a *Zygomaturus* sp. mandible found in these sediments at Batesford in 1975 (Rich, 1976). This unit has yielded a substantial list of fossilized remains known as the Dog Rocks Local Fauna, including marsupials and other mammals such as Rodentia (Muridae), along with representatives of Osteichthyes (Teleostei), Amphibia (Anura), Reptilia (Squamata), and Aves (Rich, 1991).

The sediments of this unit indicate a fluvial environment existed to the southeast of the Dog Rocks during the Pliocene. It is the characteristic preservation of the Dog Rocks local fauna, along with information

provided on the location of the new specimen by Robertson that led T. H. Rich to his assessment that the specimen came from this setting. The fossiliferous unit of the Moorabool Viaduct Sands was overlain by a number of basalt flows of the Newer Volcanics (Bowler, 1963). Three kilometres to the south of Batesford Quarry, a magnetically reversed flow with similar petrographic characteristics was dated at  $2.03 \pm 0.13$  my BP in three different samples (Aziz-ur-Rahman & McDougall, 1972). A determination for the maximum age of the fluvial sediments was provided by the foraminiferan *Globorotalia crassiformis*, which first appeared at 4.0 myBP (Whitclaw, 1989) and occurs in the sediments of the Moorabool Viaduct Sand stratigraphically below the fissures. In these sediments, the magnetic orientation is reversed (Whitclaw, 1989) which suggests that the age of the fissures is within the interval of 4.0 and 1.9 myBP, thus placing it in either the early Matuyama Chron (2.48-2.03 myBP) or the late Gilbert Chron (4.88-3.40 myBP) (Aziz-ur-Rahman & McDougall, 1972). The presence of rodents in the local fauna such as the two identified *Pseudomys* species suggests that the early Matuyama Chron is the most likely interval represented in these sediments (Whitclaw, 1989).

#### SYSTEMATIC PALAEOONTOLOGY

Subclass MARSUPIALIA Illiger, 1811

Cohort AUSTRALIDELPHIA Szalay, 1982

Order DASYUROMORPHIA Gill, 1872

Family DASYURIDAE Goldfuss, 1820

Genus *Glaucodon* Stirton, 1857

Type species *Glaucodon ballaratensis* Stirton, 1857

*Holotype*. The diagnostic features of the holotype of *G. ballaratensis* (NMV P 16136) were defined as the following: Molar row length intermediate between *Dasyurus maculatus* and *Sarcophilus laniarius harrisi*; gross morphology of mandible less robust than *Sarcophilus* form but more so than *D. maculatus*; M4 length of 10.0 mm  $\pm$  0.1 mm; M4 width 6.7 mm  $\pm$  0.4 mm; M1 length 7.2 mm  $\pm$  0.5 mm; M1 width 4.3 mm  $\pm$  0.2 mm (Stirton, 1957).

The dental morphology of the holotype of *G. ballaratensis* (NMV P16136) has been discussed by

Gill (1953), Stirton (1957), Ride (1964), Marshall (1973), Archer (1976, 1982), Areher and Bartholomai (1978), Crabb (1982) and Long et al. (2002). It was fully described by Stirton (1957) based on a solitary right mandible with M1 and M4 present, and alveoli for M2, M3, P1, P2, C, and an incomplete row of incisor alveoli. Gill (1953) postulated in his field notes, but did not formally describe, the specimen as a member of the genus *Sarcophilus*. *G. ballaratensis* is regarded herein as a morphologically intermediate form between *Sarcophilus lamarius harrisii* and *Dasyurus maculatus*, displaying characteristics referable to both species. Stirton (1957) also noted some dental morphological similarities between *G. ballaratensis* and *Dasyurus viverrinus* (Shaw, 1800), however the similarities were less marked than those of *D. maculatus*.

***Glaucodon ballaratensis* Stirton, 1957**

Fig. 1, 2, 3


*Sarcophilus*? Gill, 1953, p. 87

*Glaucodon ballaratensis* Stirton, 1957, p. 129-133, text-figure 6.

*Glaucodon ballaratensis* Stirton, 1957, - Areher, 1984, p. 634, Table 1; p. 635, fig 2; p. 639, fig 17; p. 640, fig. 18; . Pl. 6.7-1. - Crabb, 1982, p. 514 - 515, fig 4, - Areher, 1982, p. 409, fig 7, p. 417. - Ride, 1964, p. 109, fig 8; Long et al., 2002, p. 55, fig p. 53 (mislabelled as *Dasyurus dunnalli*)

*New material.* NMV P207018. As for *G. ballaratensis* holotype (NMV P16136), with the following additional features: Mandible with four molars (M1, M2, M3 and M4) in situ, however the protoconid of M4 is absent, possibly due to

Fig. 1. NMVP 207018, occlusal view. Scale bar intervals represent 10mm.



postmortem abrasion. Also present are two well-spaced premolars (P2 and P3) and a large crescentic canine. Occlusal surfaces are relatively unworn, suggesting that the specimen was a young adult. The mandible of the specimen is somewhat shorter and more robust than holotype.

*Locality of new material.* Specimen found by P. Robertson in the Moorabool Viaduct Sand, a member of the Brighton Group (Pliocene), at Portland Cement Limited's limestone quarry at Batesford, Geelong, Victoria (38° 6.5' S., 144° 17.5' E).

*Diagnosis of species.* Diagnostic features are: deep, channel-like precingulum extending to the parastylid on M2, M3 and M4 (precingulum on M1 present but less markedly) consistent with 'bone cracking' dasyurids; more robust trigonids than holotype (M1 trigonid longer, M4 trigonid both longer and wider); crowding of molar row out of line comparable to holotype (and *S. moornaensis*, but significantly less than *S. lanarius harrisii*); talonid-like notch in ante-

rior base of P2 undercuts and is rectilinear to the protoconid, premolar diastema intermediate between *D. maculatus* and *S. lanarius harrisii* and referable to alveoli on *G. ballaratensis* holotype; P1 significantly lower than P2 with rounded, low cusp. The mandible is more robust and deeper than holotype, deepest beneath the M4 between roots.

Measurements were taken from twenty specimens each of *D. maculatus* and *S. lanarius harrisii*, as well as the holotypes of *S. moornaensis* (NMV P28684) and *G. ballaratensis* (NMV P16136) to provide a morphometric comparison of twenty-eight dental and mandibular characteristics against the new specimen (Table 1). Measurements from the canine are discarded as being diagnostic because of wear via occlusion. The parameters of variability of the measurements of the talonid and trigonid basins of the M2 and M3 are not used as a diagnostic feature from the holotype, as these teeth are represented by alveoli only.

NMV P207018 differs from the holotype of *G. ballaratensis* in the following characteristics: more robust mandible, mandibular depth beneath P2 deeper

Fig. 2. NMVP 207018, lingual view. Scale bar intervals represent 10mm.





and closer to the mandibular depth proportions displayed in *S. moornaensis*, M1 and M4 wider.

NMV P207018 shares the following features with the holotype: molar talonids weaker and more adpressed than *Sarcophilus*, but less so than *Dasyurus*, molar protoconids high and sharp, particularly on M3 and inferred on M4, molar row length and crowding intermediate between *S. moornaensis* and *D. maculatus*.

It has been suggested that Robertson's specimen may represent an ontogenetically older individual of *G. ballaratensis* (G. D. Sanson, pers. comm., 1999). Whilst this addresses the apparent crowding of the molar row of the illustrated specimen (as a result of interstitial wear), the occlusal wear patterns on the protoconid and the hypoconid of the M1 of NMV P207018, unknown to Sanson, suggest it was of similar or younger ontogenetic age than the holotype. The observed differences in dental morphology between the two specimens, however, provide insufficient evidence for the existence of two species of *Glaucodon*.

Morphometric analyses set forth by Simpson, Roe

& Lewontin (1960) were used by Marshall (1973) to synonymise *G. ballaratensis* with a specimen from the Moorna Sands, South Australia. This diagnosis was later re-examined by Crabb (1982) using morphological observations, and the Moorna Sands specimen was erected as the holotype of *Sarcophilus moornaensis*. This suggests that Simpson's form of analysis is problematic when morphological observations are not included in the statistical analysis. It is pertinent to note that data derived from Simpson's approach implied the presence of two species of *Glaucodon* on the basis of dental measurements (Table 2) when comparing the *Glaucodon* specimens.

The present authors consider that it is therefore necessary to incorporate both observations of morphology and dental measurement statistics concurrently to diagnose this species from other morphologically similar taxa.

*Description of NMV P207018.* Dentition more complete than holotype (NMV P16136), however, ramus length and length of molar row is marginally shorter,

Fig. 3. NMVP 207018, labial view. Scale bar intervals represent 10mm.



CHARACTER	<i>Dasyurus maculatus</i> (n) = 20	<i>Sarcophilus lauriarius harrisi</i> (n) = 20	<i>Sarcophilus moornensis</i> (n) = 1	<i>G. ballaratensis</i> holotype MV P16136 (n) = 1	<i>G. ballaratensis</i> MV P702018 (n) = 1
Length of molar series	27.7 +/- 3.8	44 +/- 4.7	38.5	36.4	35
M1 length	7.7 +/- 2.1	9.2 +/- 2.6	7.8	7.4	6.5
M1 width	3.7 +/- 1	6.2 +/- 2.5	5.2	4	4.5
M1 trigonid length	2.7 +/- 1.2	7 +/- 3.2	7.2	4.3	4.7
M1 trigonid width	1.8 +/- 0.8	4.5 +/- 1.7	5	4.3	4.3
M1 talonid width	2.1 +/- 1	4.9 +/- 2.2	3.2	4.5	4.3
M2 length	6.5 +/- 1.7	10.9 +/- 2.9	9.5	absent	8.6
M2 width	3.4 +/- 1	6.5 +/- 1.8	5.9	absent	4.8
M2 trigonid length	3.3 +/- 1.3	8.4 +/- 2	8.7	absent	6.1
M2 trigonid width	3.4 +/- 1.2	5.2 +/- 3.7	3.8	absent	4.9
M2 talonid width	3.1 +/- 1	3.3 +/- 1.8	4.5	absent	5.2
M3 length	6.7 +/- 2.2	12 +/- 2.7	10	absent	9.8
M3 width	3.9 +/- 1.9	7.4 +/- 1.5	6.3	absent	6.7
M3 trigonid length	3.6 +/- 1.8	7.7 +/- 1.8	8	absent	7.6
M3 trigonid width	3.6 +/- 1.4	5.6 +/- 1.4	5.4	absent	5.2
M3 talonid width	2.8 +/- 1.2	2.3 +/- 2.1	3.3	absent	4.4
M4 length	6.8 +/- 2.3	11.9 +/- 3.6	11.2	10	10.1
M4 width	3.7 +/- 1.6	6.5 +/- 2.1	6.5	5.8	6.3
M4 trigonid length	4.4 +/- 1.3	9.9 +/- 1.2	10.2	6	8.2
M4 trigonid width	3.7 +/- 1.2	4.8 +/- 1.4	4.6	3.3	5.6
M4 talonid width	2.9 +/- 1	2.3 +/- 1.8	2.9	2	2.1
P2 length	4.2 +/- 1.7	6 +/- 2.7	absent	absent	4.3
P2 width	3.2 +/- 1.3	6.7 +/- 2.3	absent	absent	3.5
P3 length	4.9 +/- 2.1	6.7 +/- 2	absent	absent	5.9
P3 width	2.9 +/- 1.1	6.5 +/- 2.2	absent	absent	3.9
Mandible depth below P2	10.6 +/- 2.3	12.4 +/- 3.4	14.6	10.1	12.2
Mandible depth below M4	15.1 +/- 3.1	12.3 +/- 2.8	20	17.5	17.1
Mandible depth below M1	14.1 +/- 3.8	13.8 +/- 3	14.7	13.7	13

Table 1. Mean dental measurements from subjective and comparative material



incisor alveolus not present due to incomplete preservation of specimen. Premolars in line with significant spacing, canine relatively robust and crescentic. P1 is set oblique to antero - posterior axis of tooth row and P2 is only slightly so. Molars characterized by deep, channel-like precingulum extending to the parastylid.

M1 short but wide; heavy median protoconid; anterior median crest from base to top of protoconid; metaconid tightly adpressed to posterolingual slope of protoconid; hypoconid massive and slightly crescentic; hypoconulid in posteromedian position at posterior end of hypoconid crescent; entoconid as small rounded cusp; talonid basin with narrow lingual opening anterior and posterior to entoconid - no direct posterior opening; posterior lower labial corner extends farther posteriorly than other parts of the tooth; trigonid long and deep; paraconid obscure due to massive nature of protoconid.

M2 with moderate paraconid - protoconid shear; protoconid higher than paraconid, separated by carnassial notch; protoconid with posteriorly oriented curvature, inner lingual surface flattened; metaconid much lower than protoconid but only marginally shorter than paraconid; talonid basin comparatively large, high and shallow with wider lingual opening posterior to entoconid and small posterior opening; margin of talonid with entoconid only marginally lower than hypoconid; posterior lower labial corner marginally extends further posteriorly than other parts of the tooth; trigonid basin high and shallow.

M3 with high sharp paraconid - protoconid shear; protoconid higher than paraconid, separated by carnassial notch along paracristid; protoconid high with crescentic distally oriented curve, noticeable groove along labial aspect of metacristid; metaconid lower than protoconid but roughly the same height as paraconid; protoconid higher than paraconid.

M4 wide; protoconid absent, possibly due to abrasion; remains of high paraconid - protoconid shear surface evident; partial protoconid and paraconid separated by a carnassial notch in the paracristid; metaconid lower than paraconid; trigonid basin wide and shallow. The trigonid length compared to the total length of the molar is large, and is a characteristic of this species not shown in the comparative material of *D. maculatus*, *S. harrisi* and *S. moornaensis*.

Paraconid and metaconid progressively enlarge from M2 to M4, however talonid remains adpressed in each case. Remains of the incomplete M4 protoconid suggests it was higher than paraconid with marginally adpressed talonid present.

P2 composed of single high anterior cusp with angular posterior slope to mandible attachment longer than anterior slope; anterior attachment to mandible separated by notch in base in line with single high cusp. P1 much lower than P2; cusp barely discernible on worn surface; anteriorly well spaced from P2 but crowded in line with C.

C high and crescentic; slight lingual slope; single rounded high cusp; anteriolingual surface marked by vertical groove.

*Discussion of phylogeny of Glaucodon* Considerable discussion on the phylogeny of *Glaucodon* exists. Marshall (1973, p. 156) considered that there was an absence of morphological differences between *G. ballaratensis* and *S. moornaensis*, and hence synonymised the two species. The methodologies of Simpson, Roe & Lewontin (1960) were applied as a basis of dental morphometric analysis in this instance. While there are distinct morphological similarities between *Glaucodon* and *Sarcophilus*, specimens of the latter are larger and more robust, with a significantly greater mandibular depth and crowding of the molar row.

Ride (1964) proposed that *G. ballaratensis* was structurally ancestral to *S. harrisi*, but in some characteristics closer to larger species of *Dasyurus* such as *D. maculatus*. This suggestion implies *Glaucodon* is ancestral to *Sarcophilus* with some characteristics referable to *Dasyurus*. The phylogenetic relationships of *Dasyurus* and *Sarcophilus* have been established by molecular studies indicating that the genera form a monophyletic group (Krajewski et al., 1994). It is noteworthy that more recent molecular data is not in accord with this interpretation (Krajewski et al., 2000), but more recent morphology-based numerical parsimony supports the case for monophyly of this taxa (Wroe et al., 2000). This demonstrates the significance of morphology in the elucidation of phylogeny in extant taxa. Archer and Bartholomai (1978, p. 5) recognized "a structural lineage starting with a hypothetical fossil form similar to *D. maculatus*, to *G. ballaratensis*, to the annectant extinct species of *Sarcophilus* and ending with *S. harrisi* is one of the best documented marsupial structural phylogenies", where the annectant species mentioned is *S. moornaensis*. Wroe (1998) discussed the phylogeny of this lineage from the perspective of an ancestral Miocene dasyurid, *Ganbulanyi djadjingnli* Wroe, 1998, noting synapomorphies of *Dasyurus*, *Glaucodon* and *Sarcophilus*.

Crabb (1982) considered the lower molars of *S.*



moornaensis similar to (and as apomorphic as) the molars of *G. ballaratensis*. The mandibular depth and crowding of the molar row of *S. moornaensis* is greater than that of both specimens of *Glaucodon*, however this characteristic is less marked in the Batesford specimen. The more robust trigonids of the M1 and M4 and the mandibular depth of the Batesford specimen differ from the *G. ballaratensis* holotype, however these differences are regarded as being of infraspecific importance. It is important to consider, when discussing the relationships of the *Glaucodon*-*Dasyurus*-*Sarcophilus* lineage, the relative ages of all the taxa involved. *S. moornaensis* is considered Early Pleistocene in age (Long et al., 2002). The revised estimation of the geological age of the holotype of *G. ballaratensis* (2.0 myBP  $\pm$  0.1 myBP) removes the importance of the specimen in 'stage-of-evolution' correlation for this group of dasyurids, as the occurrence of the species is too recent to constitute 'common ancestry' (Turnbull, Lundelius & Tedford, 1993). However, the morphology of the teeth attributed herein to *G. ballaratensis* combined with the estimated geological age of NMV P207018 (2.2 myBP  $\pm$  0.25 myBP) raises the significance on the species in terms of 'stage-of-evolution' correlation. *G. ballaratensis* can now be regarded as representing a collateral lineage to *Sarcophilus*. The retention of characteristics such as the gracile mandibular form shared by *Glaucodon* and *Dasyurus* are considered symplesiomorphic.

The dental and mandibular characteristics of NMV P207018 provide new data on *Glaucodon* and the characteristics that it shares with *Sarcophilus*. *G. ballaratensis* is a Pliocene form that, whilst sufficiently different to warrant generic distinction from

the Dasyurinae and *Sarcophilus*, has characteristics ally-ing it with these taxa. The molar row crowding and proportions of the M4 trigonid of NMV P207018 implies a closer affinity between the genera *Glaucodon* and *Sarcophilus* than previously postulated. The poorly developed molar talonids of NMV P207018 align the specimen with *Glaucodon* rather than *Sarcophilus*. The robust molars of the Batesford specimen are morphologically intermediate between the holotype of *G. ballaratensis* and *S. moornaensis*. However, these differences are likely to represent intraspecific variation within *G. ballaratensis* rather than a new species of *Glaucodon*.

A possible interpretation of the phylogeny of the lineage is as follows: at some point prior to the late Pliocene, the Dasyurinae diversified resulting in the origin of the *Sarcophilus*-*Glaucodon* lineage. The plesiomorphic gracile morphology and uneroded molars of the ancestral form were retained and are present in extant species of *Dasyurus*. The *Sarcophilus*-*Glaucodon* lineage retained some elements of the dental arrangement of the ancestral form. However, significant progressive crowding of the molar row is an evident apomorphy of the subfamily as displayed in *S. lanarius harrisi*.

*Glaucodon* branched from the *Sarcophilus*-*Glaucodon* lineage after the split from the Dasyurinae, but prior to the occurrence of *S. moornaensis*. *G. ballaratensis* arose at some point prior to the Late Pliocene, and retained the symplesiomorphic proportions present in the mandible of *D. maculatus*, but also displayed characteristics which unite it with *Sarcophilus*. A characteristic of *Sarcophilus* is the gradual increase of morphological robustness through the Plio-Pleistocene *S. moornaensis* to the massively

CHARACTER	<i>G. ballaratensis</i> holotype MV P 16136	<i>G. ballaratensis</i> MV P 702018	Log. difference	Nearest whole number value for V	Indicative of new species ?
Length of molar series	34.4	35	0.0076	<1	N
M1 Length	7.4	6.5	0.056	3 to 4	Y
M1 Width	4	4.5	0.0511	3 to 4	Y
M1 Trigonid length	4.4	4.7	0.029	1 to 2	N
M1 Trigonid width	4.3	4.3	nil	N/A	N
M1 Talonid width	4.5	4.3	0.03	1 to 2	N
M4 Length	10	10.1	0.004	<1	N
M4 Width	5.8	6.3	0.0359	2 to 3	N
M4 Trigonid length	6	8.2	0.1356	7 to 8	Y
M4 Trigonid width	3.3	5.6	0.2297	12 to 13	Y
M4 Talonid width	2	2.1	0.0212	1 to 2	N
Mandible depth below P2	10.1	12.2	0.082	4 to 5	Y
Mandible depth below M4	17.3	17.1	0.0051	<1	N
Mandible depth below M1	13.5	13.2	0.0097	<1	N

Table 2. Application of morphometric analysis of Simpson et al. (1960) to *Glaucodon*. (Note: values for V greater than 3 indicate a different species. Chronology and phylogeny are not applied in this approach to species differentiation.)

robust *S. lanianus* complex of the Pleistocene. Future discoveries from these theorised lineages could add to the understanding of the evolution of the Dasyuridae in the Plio-Pleistocene.

#### ACKNOWLEDGEMENTS

The authors are grateful to Deakin University for funding and facilities, Dr. T. H. Rich, and Mr. P. R. Brown provided valuable comments during the course of this study. Museum Victoria arranged specimen loans and access to Museum Victoria collections. Assoc. Prof. G. D. Sanson of Monash University, N. W. Longmore of Museum Victoria and Danielle O'Neill provided useful advice in constructing this paper.

#### REFERENCES

- ARCHER, M., 1976. The dasyurid dentition and its relationships to that of didelphids, thylacynids, borhyaenids (Marsupicarnivora) and peremylids (Peremylina: Marsupialia). *Australian Journal of Zoology, Supplementary Series*, 39: 1-34.
- ARCHER, M., 1982. Review of the Dasyurid (Marsupialia) fossil record, integration of data bearing on phylogenetic interpretation, and suprageneric classification. In *Carnivorous Marsupials Vol. 2*, Archer, M., ed. Royal Zoological Society of New South Wales, Sydney.
- ARCHER, M., & CLAYTON, G. (eds) 1984. *Vertebrate Zoogeography & Evolution in Australasia (Animals in Space and Time)*, Hesperian Press, Victoria Park.
- ARCHER, M. & BARTHOLOMAI, A., 1978. Tertiary mammals of Australia: A synoptic view. *Alcheringa* 2: 1-20.
- AZIZ-UR-RAHMAN, & McDUGALL, I., 1972. Potassium-Argon ages on the New Volcanics of Victoria. *Proceedings of the Royal Society of Victoria* 85: 61-9.
- BOITARD, P., 1842. *Le Jardin des Plantes a Paris. Description et moeurs des Mammiferes de la Menagerie et du Museum d'Histoire Naturelle/ precede d'une introduction historique, descriptive et pittoresque par M. J. Janin*, Paris: J.J. Dubochet et Ce, i-lxvi, 472 pp.
- BOWLER, J. M., 1963. Tertiary Stratigraphy and Sedimentation in the Geelong - Maude area, Victoria. *Proceedings of the Royal Society of Victoria* 76: 7-137, Pls. 15-18.
- CRABB, P. L., 1982. Pleistocene dasyurids (Marsupialia) from south western New South Wales. In *Carnivorous Marsupials. Vol. 2*, M. Archer, M., ed. Royal Zoological Society of New South Wales, Sydney. 511-516.
- GERDTZ, W. R., 2001. McCoy and *Sarcophilus harrisii* Boitard, 1842 - a Diabolical Relationship. *The Victorian Naturalist* 118 (5), 231-233.
- GILL, E. D., 1953. Distribution of the Tasmanian Devil, the Tasmanian Wolf and the Dingo in southeast Australia in Quaternary time. *The Victorian Naturalist* 70: 86-90.
- GILL, T., 1872. Arrangement of the Families of Mammals: with analytical tables. Prepared for the Smithsonian Institution by Theodore Gill. *Smithsonian miscellaneous collections*; Volume 9, article 1, number 230, 1-98.
- GOLDFUSS, G. A., 1820. Thl. 3 Handbuch der Zoologie. In *Handbuch der Naturgeschichte: zum Gebrauch bei Vorlesungen*. G. H. von Schubert, Nuremberg.
- ILLIGER, J. K. W., 1811. C. Illigeri ...*Prodromus systematis mammalium et avium additis terminis zoographicis utriusque classis, eorumque versione Germanica*. Berolini.
- KERR, R., 1792. *The Animal Kingdom, or zoological system, of the celebrated Sir Charles Linnaeus; Class I. Mammalia, containing a complete systematic description, arrangement, and nomenclature, of all the known species and varieties of the Mammalia, or animals which give suck to their young; being a translation of that part of the Systema Naturae, as lately published, with great improvements, by Professor Gmelin of Goettingen. Together with numerous additions from more recent zoological writers, and illustrated with copper plates*. London: J. Murray & R. Faulder, Part 1, xi + 400 pp. 9 pls.
- KRAJEWSKI, C., PAINTER, J., BUCKLEY, L. & WESTERMAN, M., 1994. Phylogenetic structure of the marsupial family Dasyuridae based on Cytochrome - DNA sequences. *Journal of Mammalian Evolution* 2 25-35.
- KRAJEWSKI, C., WROE, S. & WESTERMAN, M., 2000. Molecular evidence for the pattern and timing of cladogenesis in dasyurid marsupials. *Zoological Journal of the Linnean Society*, 130: 375-404.
- LONG, J., ARCHER, M., FLANNERY, T. & HAND, S.,



2002. *Prehistoric Mammals of Australia and New Guinea: One Hundred Million Years of Evolution*. University of New South Wales Press Ltd, University of New South Wales, Sydney.
- MARSHALL, L. G., 1973. Fossil vertebrate faunas from the Lake Victoria region, S. W. New South Wales, Australia. *Memoirs of the National Museum of Victoria* 34: 151-81.
- OWEN, R. In T. L. Mitchell, 1838. *Three expeditions into the interior of eastern Australia, with descriptions of the recently explored region of Australia Felix, and of the present colony of New South Wales*. T & W Boone, London.
- OWEN, R., 1877. *Researches on the fossil remains of the extinct mammals of Australia; with a notice on the extinct marsupials of England*, Volumes 1 & 2. J. Erxleben, London.
- RICH, T. H., 1976. Recent fossil discoveries in Victoria. *Victorian Naturalist* 93(5): 198-206.
- RICH, T. H., 1991. Australia's mammal record. In *Vertebrate Palaeontology of Australasia*, P. Vickers-Rich, J.M. Monaghan, R. F. Baird, & T. H. Rich, eds. Monash University Publishing Committee, Melbourne, 893-1070.
- RIDE, W. D. L., 1964. A review of the Australian fossil marsupials. *Journal of the Proceedings of the Royal Society of Western Australia*. 47: 97-131.
- SHAW, G., 1800. *General Zoology or systematic natural history. Quadrupeds*. G. Kearsley, London, Vol. 1, Pt 2, i-viii, 249-552.
- SIMPSON, G. G., ROE, A. & LEWONTIN, R. C., 1960. *Quantitative Zoology*. Harcourt, Brace & World Publishing Inc., New York.
- STIRTON, R. A. 1957. Tertiary Marsupials from Victoria, Australia. *Memoirs of the National Museum of Victoria* 21: 121-134.
- SZALAY, F. S., 1982. A new appraisal of marsupial phylogeny and classification. In *Carnivorous Marsupials Volume 2*. M. Archer ed., Surrey Beatty & Sons Pty. Ltd. and The Royal Zoological Society of New South Wales, Mosman, 621-640.
- TURNBULL, W. D. & LUNDELIUS, E. L. fil., 1970. The Hamilton Fauna: A late Pliocene mammalian fauna from the Grange Burn, Victoria, Australia. *Fieldiana Geology* 19: 1-163.
- TURNBULL, W. D., LUNDELIUS, E. L. fil. & TEDFORD, R. H., 1993. Fossil vertebrate locality at Smeaton, Victoria. *Memoirs of the Association of Australasian Palaeontologists* 15: 429-440.
- WHITELAW, M. J., 1989. Magnetic polarity stratigraphy and Mammalian fauna of the late Pliocene (Early Matuyama) section at Batesford, Victoria. *Journal of Geology* 97: 11-24.
- WROE, S., 1998. A new 'bone-cracking' dasyurid (Marsupialia) from the Miocene of Riversleigh, northwestern Queensland. *Alcheringa* 22: 277-284.
- WROE, S., 2003. Australian marsupial carnivores: an overview in recent advances in palaeontology. In M. Jones, C. Dickman, & M. Archer, eds. *Predators with Pouches: The biology of Carnivorous Marsupials*, CSIRO Publishing, Melbourne, 102-123.
- WROE, S., EBACH, M., AIYONG, S., MUIZON, C. de, & MUIRHEAD, J., 2000. Cladistic analysis of dasyuromorphian (Marsupialia) phylogeny using cranial and dental features. *Journal of Mammalogy*, 81: 1008-1024.
- WROE, S., & MACKNESS, B. S., 2000. A new genus and species of dasyurid from the Pliocene Chinchilla Local Fauna of south-eastern Queensland. *Alcheringa* 24: 319-326.

Manuscript received 16 June 2003

Revision accepted 17 November 2003



AN EARLY OCCURRENCE OF *SARCOPHILUS LANIARIUS HARRISII*  
(MARSUPIALIA, DASYURIDAE) FROM THE EARLY PLEISTOCENE OF  
NELSON BAY, VICTORIA.

W. R. GERDTZ & N. W. ARCHBOLD

School of Ecology and Environment, Melbourne Campus, Deakin University, Burwood, Victoria, 3125

GERDTZ, W. R. & ARCHBOLD, N. W., 2003:11:30. An early occurrence of *Sarcophilus laniarius harrisii* (Marsupialia, Dasyuridae) from Nelson Bay, Victoria. *Proceedings of the Royal Society of Victoria* 115(2): 45-54. ISSN 0035-9211.

The isolated lower molar series of a dasyurid from early Pleistocene sediments at Nelson Bay, near Portland, is referred herein to *Sarcophilus laniarius harrisii* (Boitard, 1842). Dental measurements and morphological comparisons taken from this tooth series compare closely with those of the extant subspecies. The genus *Sarcophilus* is known from the early Pleistocene by the species *S. moornaensis* Crabb 1982. Fossil material assigned to *S. laniarius harrisii* is now also known from Early Pleistocene sediments. The new specimens from Nelson Bay extend the origin of *S. laniarius harrisii* to the early Pleistocene.

*Key words:* *Sarcophilus laniarius harrisii*, Pleistocene, Portland, Nelson Bay Formation, Dasyuromorphia, Dasyuridae.

THE FIRST reports of the occurrence of the fossil remains of the living Tasmanian *Sarcophilus* in the Quaternary deposits of Victoria were by Selwyn (1858, 1860) based on identifications by Frederick McCoy. The modern geographical distribution of the extant dasyurid subspecies *Sarcophilus laniarius harrisii* (Boitard, 1842) widely known as *S. harrisii* (Boitard 1842) is restricted to dry sclerophyll forests and coastal woodlands of Tasmania (Strahan, 1995). The fossil record of the species has a distribution on mainland Australia during the Pleistocene (Long et al. 2002). Fossil *S. laniarius* has been recorded in many geochronologically dated Pleistocene and early Holocene terrestrial and cave fossil localities. Representative occurrences are provided in Table 1. The maximum range of the geochronologically dated specimens is no older than 0.352 +/- 1.8 Mya down to less than 10,000 years. The extinction of the species from the mainland is believed to have been the result of a failure of the species to compete successfully with the introduced Dingo *Canis familiaris dingo* (see Corbett, 1985), which occupied a comparable ecological niche (Strahan, 1995).

The earliest occurrence of *Sarcophilus* is from the early Pleistocene, with *S. moornaensis* Crabb (1982), known from a right mandible and molars, and is a member of the Fisherman's Cliff Local Fauna (Crabb, 1982). Crabb's species comes from the sands of the Moorna Formation in south western New South Wales, which is estimated to be early Pleistocene in

age (Long et al. 2002). Late Pleistocene fossil-bearing deposits in the Wellington Caves, New South Wales have yielded significantly larger specimens of *Sarcophilus*, and the species *S. laniarius* (Owen, 1838) was proposed for this material. Werderlin (1987) argued strongly that *S. laniarius* is the senior synonym for *S. harrisii*, and recognised three subspecies of *S. laniarius*, namely *S. laniarius laniarius* (Owen, 1838), *S. laniarius dixonae* (Werderlin, 1987) and *S. laniarius harrisii* (Boitard, 1842), the latter of which is also the extant subspecies. Dawson (1982) recognised the basis for diagnostic distinction of *S. laniarius laniarius* from the extant species is the larger size rather than discrete morphological differences. Murray (1991) noted that the Wellington Caves *S. laniarius laniarius* is an "essentially statistically assigned (sub) species (with) no particular morphological features other than significantly larger size and associated allometry from ...present-day Tasmanian Devils" (Murray, 1991, p. 1090-1091). As a result we accept that *S. laniarius laniarius* is a larger form of *S. laniarius harrisii*. For the purposes of this present report, we follow Werderlin's (1987) taxonomy of the three subspecies, and the binomen *S. laniarius* (Owen, 1838) has priority, as shown in Long et al. (2002).

It may be argued that the use of *S. laniarius* (Owen, 1838) as the senior synonym of *S. harrisii* (Boitard, 1841) for the living Tasmanian Devil results in the destabilization of commonly used nomenclature. While Environment Australia's biodiversity fauna list

(Environment Australia, Australian Biological Resources Study, 2002) refers to *S. harrisii* as the binomen for the living species, it is in contrast with Smithsonian Museum of Natural History taxonomic database (Smithsonian Museum of Natural History, Department of Systematic Biology, 1993) that recognizes the priority of *S. lamarius* for the extant species. As shown by Werderlin (1987) and subsequent workers (Long *et al.* 2002), this recognition of priority does not significantly destabilize nomenclature and is therefore retained by us.

In the late 1970's, a stratigraphical review of the sediments outcropping in Nelson Bay, near Portland, was undertaken by E. D. Gill, C. W. Mallett, J. M. Lindsay and N. W. Archbold. Archbold collected a small collection of fossil bone and tooth material from a single locality. Amongst the collected material were assorted diprotodontid, vombatid, macropodine and dasyurid mandibular and dental remains. It is the dasyurid material that is described herein and is assigned to *S. lamarius harrisii* on the basis of dental measurements and morphology. Due to the age of the sediments that contained this material (0.73 - 1.66mya) (White, 2002), it appears that they constitute the earliest occurrence of this species.

#### *Stratigraphy and age*

Based on field notes and personal recollections from one of us (N. W. A.) relating to the locality, the unit in which the material is referred to is unquestionably the Nelson Bay Formation, located approximately 5 km south of Portland (38° 36' S., 141° 35' E.). Studies of the sedimentary processes and the formation of the bone beds which outcrop at Nelson Bay have been undertaken by Boutakoff (1963) and Kenley (1988), and were reviewed by White (2002).

The sediments of the Nelson Bay Formation were deposited in a lacustrine environment that existed at this locality at the time of deposition. The lake occupied a large subsided caldera formed from volcanic activity 2.76 million years ago (Aziz-Ur-Rhamen & McDougal, 1972). Sediments from the surrounding Portland Limestone were levelled and redeposited as sedimentary beds within the 'Nelson Bay caldera lake' (White, 2002). These sediments in turn formed calcarenite beds, interspersed with clay beds. The clay beds in the caldera lake sediments are marked by the presence of rip-up clasts formed by strong currents during periods of flooding of the lake,

and form a portion of the Nelson Bay Formation.

The Nelson Bay Formation includes a distinct unconformity, forming a boundary between the Upper Nelson Bay Formation and the Lower Nelson Bay Formation. This unconformity has been correlated with the Gunz Glacial Peak (Boutakoff, 1963), implying a minimum age of 0.73my for the underlying Lower Nelson Bay Formation. The sediments of the Lower Nelson Bay Formation have been described as approximately 10m of clay beds, which includes approximately 3m of horizontally bedded calcarenite, resting on 2m of sandy clay. This in turn rests upon pink sandy clay soils (Boutakoff, 1963). It is the sandy clay soils that form the fossil bone beds in the Lower Nelson Bay Formation. The presence of the foraminiferal species *Globorotalia truncatulinoides* in the sediments of the Nelson Bay Formation implies a maximum age of 1.9my for this unit (Srinivasan & Kennet, 1981). Magnetic polarity stratigraphy studies have concluded that the Nelson Bay Formation represents an age of 1.63 - 0.71 my (Whitelaw, 1991). MacFadden (1987) noted that the Nelson Bay Formation is significant in terms of palaeontological, radio isotopic and palaeomagnetic data leading to a relatively precise geochronology for the formation (MacFadden *et al.* 1987). Kenley assigned the Nelson Bay Formation to the Early Pleistocene (Kenley 1988, p. 366). Rich *et al.* (1991) also stressed that the Nelson Bay Local Fauna "is the single securely dated Early Pleistocene assemblage from either Australia or Papua New Guinea." (p. 1014). White (2002) noted that five distinct fossil bone beds are present in the Lower Nelson Bay Formation. The collection of the material described here can not be located precisely to a particular bone bed described, but it originated from the Lower Nelson Bay Formation approximately 50m west of the sewer outfall, and was at the time of collection approximately 1.5m above the beach sand in front of the cliff. The material can therefore be considered Early Pleistocene in age.

#### *The Nelson Bay Local Fauna*

Hann (1983) partially described the 'Nelson Bay Local Fauna' in an unpublished Honours thesis, which was subsequently updated by Herrmann (2000) in an unpublished Masters thesis. Whitelaw (1991) referred to the Nelson Bay Local Fauna in terms of geochronology, however the composition of this Local Fauna was not included in this reference. White (2002) compiled a list of identified fossil species from this



locality which comprise the fauna identified prior to 2000 and included additional material in the local fauna which was identified for the purposes of that study. A combined lists of species from the Nelson Bay Local Fauna is provided in Table 2.

#### SYSTEMATIC PALAEONTOLOGY

Subclass MARSUPIALIA Illiger, 1811

Cohort AUSTRALIDELPHIA Szalay, 1982

Order DASYUROMORPHIA Gill, 1872

Superfamily DASYUROIDEA Goldfuss, 1820

Family DASYURIDAE Goldfuss, 1820

Genus *Sarcophilus* Geoffroy Saint-Hilaire & Cuvier, 1837

Type Species *Sarcophilus laniarius* (Owen, 1838)

#### Selected synonymy.

*Didelphis ursina* Harris 1808: 176, pl. 19.

*Dasyurus laniarius* Owen 1838: 369, pl. 49, figs 3-7.

*Sarcophilus harrisii* Boitard 1842: 290, Merigian *et al.* 2002: 84, fig. 2C, fig 8A-F.

*Sarcophilus (Dasyurus) ursinus* Owen, 1877: 105, pl. 13, figs 2, pl. 15, fig 2.

*Sarcophilus (Dasyurus) laniarius* Owen, 1877: 105, pl. 5, figs 1-6, pl. 14, fig 2.

*Sarcophilus satanicus* Thomas, 1903: 289.

*Sarcophilus laniarius laniarius* Owen, 1838, -Werderlin, 1987: 9.

*Sarcophilus laniarius* Owen, 1838, Long *et al.* 2002: 55.

**Description of material.** The illustrated material collected at Nelson Bay comprises a complete lower molar series (M/1, M/2, M/3 and M/4), a single lower premolar (P/2), and canine. The material is composed of teeth only, traces of mandibular material were found associated with the suite of teeth but disintegrated on collection. There is no doubt that all the teeth derived from a solitary individual. The canine is crescentic, denuded of dental enamel and shares the robust morphology of the *S. laniarius harrisii* used for

Fig. 1. *Sarcophilus laniarius harrisii*, labial view. Scale bar intervals represent 10mm.





comparative purposes. The premolar (P/2) is composed of enamel and a single root consistent with *S. lanianus harrisii* morphology. M/1, M/2 and M/3 are complete teeth with enamel and roots intact, M/1 showing a degree of occlusal wear consistent with mature sarophiloid dasyurids. M/4 is fragmentary (in two portions) with much of the talonid missing, break is a vertical 'crack' from protoconid. Trigonid of M4/ present, talonid is absent.

A list of 22 morphometric characteristics of the lower dentition of *S. lanianus harrisii* were recorded from 20 individuals. This morphometric data was then compared to corresponding characteristics on the Nelson Bay material, and is shown in Table 3. The Nelson Bay material is unquestionably from *S. lanianus harrisii* on the basis of this comparison.

*Sarcophilus lanianus harrisii* Boitard, 1842

**Discussion.** Considerable discussion on the dental morphology and fossil record of *S. lanianus harrisii* is available. Archer (1976) discussed dasyurid dental nomenclature and demonstrated the relationships of

dasyurids (including *S. harrisii*) to didelphids, thylacineids, borhyaenids and peramelids. Archer (1982) reviewed the dasyurid fossil record (including *S. harrisii*), and investigated the phylogeny of dasyurids generally. Wroe *et al.* (2000) investigated the dental and cranial characteristics of dasyurids (including *S. [lanianus] harrisii*) using cladistic analysis for phylogenetic analysis. The dental morphological characteristics assigned to *S. lanianus harrisii* in these investigations conform closely to the Nelson Bay material. The illustrated specimen was also compared with the holotype of *S. moornensis* and was shown to share broad morphological affinities consistent with the generic distinction of *Sarcophilus*, but was found to differ in a combination of features, such as the generally smaller size of *S. moornensis*. The *S. moornensis* specimens also displayed more elongate M/3 and M/4 talonids, more distinct talonid cusps, better developed metaconids and more laterally compressed trigonids than the material described herein. Both the comparative *S. lanianus harrisii* material and Nelson Bay specimens have weaker metaconids than *S. moornensis* and are larger than *S. moornensis*. The new teeth are therefore assigned

Fig. 2. *Sarcophilus lanianus harrisii*, lingual view. Scale bar intervals represent 10mm.



to *S. lanarius harrisii* on the basis of close affinities to the comparative material.

Long et al. (2002), Gill (1953) and Archer *et al.*, (1984) considered the time range of *Sarcophilus* to be early Pleistocene to present day, noting the occurrence of *S. lanarius harrisii* in many Pleistocene faunas throughout mainland Australia and Tasmania. Dawson (1982), in reviewing the taxonomic status of specimens from eastern Australian fossil localities, included *S. lanarius* fossil material no older than 30,000 years old, and considered the time range of *S. lanarius* to be Pleistocene. Güler (1982) discussed the temporal distribution of *S. lanarius harrisii* on mainland Australia, citing fossil material no older than 5000 years old. Due to the geochronology of the bone beds in the Lower Nelson Bay Formation, the Nelson Bay material constitutes the geologically earliest recorded occurrence of verifiable age of *S. lanarius harrisii*.

The presence of *S. lanarius harrisii* in earliest Pleistocene sediments provides further information on the evolution of the sarcophilines. Archer and Bartholomai (1978, p. 5) recognized "A structural lineage starting with a hypothetical fossil form similar to *Dasyurus maculatus* Kerr, 1792, to *Glauconodon ballaratensis* Stirton, 1957, to the annectant extinct species of *Sarcophilus* and ending with *S. harrisii* is one of the best documented marsupial structural phylogenies", where the annectant species mentioned is *S. moornaensis*. *S. moornaensis* is considered early Pleistocene in age (Long et al. 2002). The geochronology of the sediments containing the illustrated material implies *S. lanarius harrisii* may have existed contemporaneously with *S. moornaensis*, and as such *S. moornaensis* is not considered ancestral

to *S. lanarius harrisii*. *Glauconodon ballaratensis* represents a sister taxon relationship to *Sarcophilus* and *Dasyurus*. Elsewhere in this issue the morphological similarities of *G. ballaratensis* to *Dasyurus maculatus* are discussed. The age of the illustrated material implies that the genus *Sarcophilus* arose at some point prior to the early Pleistocene from an ancestral form morphologically similar to *G. ballaratensis*. Future discoveries of Cenozoic and Quaternary sarcophiline material will provide further insight to the evolution of *Sarcophilus* and the placement of *Sarcophilus* in dasyurid phylogeny.

The new material in the Lower Nelson Bay Formation also provides additional information on the composition of the Nelson Bay Local Fauna and the implied environmental conditions of Nelson Bay in the early Pleistocene. Notably, the Nelson Bay Local Fauna is composed of an extensive and diverse list of macropodids of both the browsing and grazing dental grades. The presence of these forms implies a combination of palaeoenvironments may have surrounded the Nelson Bay Caldera Lake during the Pleistocene. The inclusion of arboreal forms such as the pseudocheirids suggest a forested environment may have existed close to the site of deposition, however the presence of grazing forms such as *Macropus* suggest grassland environments may have also been nearby.

Hann (1983) described the Nelson Bay Local Fauna as a biocenosis in composition, implying the fauna once lived together prior to death and subsequent fossilisation. The Nelson Bay Local Fauna was also regarded as being derived from catastrophic mortality due to the presence of juveniles, young adults and mature adults (Hann, 1983). The

Location	Age (in years)	Reference
Strathdownie, Victoria	"Late Pliocene"	Werderlin, 1987
Devil's Lair Cave, Western Australia	35,200 +/- 1,800	Archer, 1982
Keilor, Victoria	>18,000	Marshall, 1974
Buchan, Victoria	22,980 +/- 2,000	Flood, 1973
Lancefield, Victoria	26,000 +/- 500	Gillespie <i>et al.</i> , 1978
Laura, Queensland	10,000 app.	Horton, 1977
Lake Menindee, New South Wales	18 - 26,000	Tedford, 1967
Laker Gumpung, New South Wales	>12,000	Hope, 1978
Dempsey's Lake, South Australia	>30,000	Wells, 1978
Kangaroo Island, South Australia	16,000 +/- 100	Hope et al., 1977
Queenscliff, Victoria	<10,000	Bartholomai & Marshall, 1972
Gisborne Cave, Victoria	<10,000	Gill, 1964

Table 1. Geochronologically dated fossil *S. lanarius* material from the Pleistocene and Holocene of mainland Australia

Subclass	Order	Family	Genus and species
Prototheria	Monotremata	Tachyglossidae	<i>Zaglossus robusta</i>
Metatheria	Polyprotodontia	Thylacinidae	<i>Thylacinus cynocephalus</i>
		Dasyuridae	<i>Sarcophilus</i> c.f. <i>lanianus harrisi</i>
	Diprotodonta	Thylacoleonidae	<i>Thylacoleo</i> sp.
		Ektopodontidae	<i>Darcins 'braileyi'</i>
		Pseudocheiridae	<i>Pseudocheirus</i> c.f. <i>peregrinus</i>
			<i>Pseudocheirus</i> c.f. <i>stirtoni</i>
			Giant <i>Pseudocheirus</i> sp.'
		Diprotodontidae	<i>Diprotodon</i> c.f. <i>optatum</i>
		Palorchestidae	<i>Zygomaticus trilobus</i>
			<i>Palorchestes</i> c.f. <i>parvus</i>
		Macropodidae	<i>Sthenurus</i> sp.
			<i>Protemnodon brehus</i>
			<i>Protemnodon anak</i>
			<i>Macropus</i> sp.
			<i>Macropus</i> c.f. <i>titan</i>
			<i>Baringa nelsonensis</i>
			<i>Thylogale</i> c.f. <i>billardieri</i>
			<i>Thylogale</i> sp.
			<i>Lagostrophus</i> sp.
			<i>Onychogalca</i> sp.
			<i>Setonix brachyurus</i>
			'Very small macropodid'
Eutheria	Rodentia	Potoroidae	indet.
		Muridae	<i>Pseudomys</i> c.f. <i>shortridgei</i>

Table 2. The Nelson Bay Local Fauna, adapted from White (2002) and Herrmann (2000).



CHARACTER	<i>Sarcophilus harrisi</i> (n) = 20	<i>Sarcophilus moornaensis</i> (n) = 1	Portland <i>Sarcophilus</i> (n) = 1
M1 length	9.2	7.8	9.2
M1 width	6.2	5.2	7
M1 trigonid length	7	7.2	4.3
M1 trigonid width	4.5	5	6.4
M1 talonid width	4.9	3.2	5.9
M2 length	10.9	9.5	10.9
M2 width	6.5	5.9	8
M2 trigonid length	8.4	8.7	6.6
M2 trigonid width	5.2	3.8	5.9
M2 talonid width	3.3	4.5	4.9
M3 length	12	10	14.8
M3 width	7.4	6.3	8.7
M3 trigonid length	7.7	8	9.2
M3 trigonid width	5.6	5.4	6.6
M3 talonid width	2.3	3.3	4
M4 length	11.9	11.2	absent
M4 width	6.5	6.5	7.1
M4 trigonid length	9.9	10.2	>9.4
M4 trigonid width	4.8	4.6	>6.2
M4 talonid width	2.3	2.9	absent
P3 length	6	absent	>6
P3 width	6.7	absent	>4.4

Table 3. Comparative dental measurements of the Portland *Sarcophilus laniarius harrisi*

disconformities which truncate the bone beds in the Lower Nelson Bay Formation were correlated with periods of substantial climatic change, which may have triggered the catastrophic mortality observed by Hann (White, 2002). The disconformities represent cool, dry climates which led to the accumulation of bone material on the land surface. Subsequent flooding at the beginning of the next warm, wet phase concentrated this bone material as basal lag deposits (White, 2002). Future investigations and discoveries from the Nelson Bay Local Fauna will provide a greater insight to the nature of early Pleistocene mammalian assemblages and environmental conditions on mainland south eastern Australia.

#### ACKNOWLEDGEMENTS

The authors are grateful to Deakin University for funding and use of facilities. N. W. Longmore, Collection Manager of Ornithology and Mammalogy, Museum Victoria, arranged access to Museum Victoria Mammalogy collections, and also provided useful advice in writing this paper.

#### REFERENCES

- ARCHER, M., 1976. The dasyurid dentition and its relationships to that of didelphids, thylacinids, borhyaenids (Marsupicarnivora) and peremelids (Peremelina: Marsupialia). *Australian Journal of Zoology, Supplementary Series*, 39: 1-34.
- ARCHER, M., 1982. Review of the Dasyurid (Marsupialia) fossil record, integration of data bearing on phylogenetic interpretation, and suprageneric classification. In *Carnivorous Marsupials volume 2*, M. Archer ed., Royal Zoological Society of New South Wales, Sydney.
- ARCHER, M. & BARTHOLOMAI, A., 1978. Tertiary mammals of Australia: A synoptic view. *Alcheringa* 2: 1-20.
- ARCHER, M., CLAYTON, G. & HAND, 1984. A checklist of Australasian fossil mammals. In *Vertebrate Zoogeography & Evolution in Australasia (Animals in Space and Time)*. M. Archer & G. Clayton eds., Hesperian Press, Victoria Park.
- AZIZ-UR-RAHMAN, & McDUGALL, I., 1972. Potassium-Argon ages on the New Volcanics of Victoria. *Proceedings of the Royal Society of Victoria* 85: 61-69.
- BARTHOLOMAI, A. & MARSHALL, L. G., 1972. The identity of the supposed dasyurid marsupial *Sarcophilus prior* De Vis 1883, with comments on other reported "Pliocene" occurrences of *Sarcophilus*. *Memoirs of the Queensland Museum* 16: 396-374.
- BOITARD, P., 1842. *Le Jardin des Plantes a Paris. Description et moeurs des Mammiferes de la Menagerie et du Museum d'Histoire Naturelle/ precede d'un introduction historique, descriptive et pittoresque par M. J. Janin. J.J. Dubochet et Ce*, Paris, lxvi, 472 pp.
- BOUTAKOFF, N., 1963. The Geology and Geomorphology of the Portland area. *Geological Survey of Victoria Memoir No. 22*: 52-58.
- CORBETT, L. K., 1985. Morphological comparisons of Australian and Thai dingoes: A reappraisal of dingo status, distribution and ancestry. *Proceedings of the Ecological Society of Australia* 13: 277-291.
- CRABB, P. L., 1982. Pleistocene dasyurids (Marsupialia) from south western New South Wales. In *Carnivorous Marsupials. Volume 2*. M. Archer ed., Royal Zoological Society of New South Wales, Sydney. 511-516.
- DAWSON, L., 1982. Taxonomic status of fossil Devils (*Sarcophilus*, Dasyuridae, Marsupialia) from Late Quaternary Eastern Australian Localities. In *Carnivorous Marsupials. Volume 2*. M. Archer ed., Royal Zoological Society of New South Wales, Sydney. 517-525.
- ENVIRONMENT AUSTRALIA, AUSTRALIAN BIOLOGICAL RESOURCES STUDY (ABRS), 2002. Australian Faunal Directory Database. *Environment Australia, Federal Department of the Environment and Heritage*, 2002. <http://www.ca.gov.au/biodiversity/abrs/abif/fauna/afd/mformat.html>
- FLOOD, J., 1973. Pleistocene human occupation and extinct fauna in Clogg's Cave, Buchan, South-eastern Australia. *Nature (London)* 246(5431): 303.
- GEOFFROY SAINT-HILAIRE, E., & CUVIER, G. L. C. F. D., 1837. *Histoire naturelle des Mammiferes, avec des figures originales &c.* (Livraison 70). Blaise, Paris.
- GILL, E. D., 1953. Distribution of the Tasmanian Devil, the Tasmanian Wolf and the Dingo in southeast Australia in Quaternary time. *Victorian Naturalist* 70: 86-90.



- GILL, E. D., 1964. Appendix in Mahoney, J. A., 1964. The taxonomic status of *Dayurus affinis* McCoy, 1865 (Dasyuridae) and *Hysiprymnus tristucatus* McCoy, 1865 (Macropodidae), two marsupials from a Holocene cave near Gisborne, Victoria. *Proceedings of the Royal Society of Victoria* 77: 525-533.
- GILL, T. 1872. Arrangement of the Families of Mammals: with analytical tables. Prepared for the Smithsonian Institution by Theodore Gill. *Smithsonian miscellaneous collections*. 9 (1): 1-98.
- GILLESPIE, R., HORTON, D. R., LADD, P., MACUMBER, P. G., RICH, T. H., THORNE, R. & WRIGHT, R. V. S. 1978. Lancefield swamp and the extinction of the Australian megafauna. *Science*. 200: 1044-1048.
- GOLDFUSS, G. A. 1820. Theil 3 Handbuch der Zoologie. In *Handbuch der Naturgeschichte: zum Gebrauch bei Vorlesungen*. G. H. von Schubert. Nuremberg.
- GUILIER, E. R., 1982. Temporal and spatial distribution of the Tasmanian Devil *Sarcophilus harrisii* (Dasyuridae: Marsupialia). *Papers and Proceedings of the Royal Society of Tasmania* 116: 153-163.
- HANN, L., 1983. The Vertebrate Palaeontology and Age of the Nelson Bay Formation, Portland, Victoria. *Unpublished Honours Thesis, Department of Earth Science, Monash University, Clayton, Victoria, Australia*.
- HARRIS, G. P., 1808. Description of two new species of *Didelphis* from Van Diemen's Land. *Transactions of the Linnaean Society, London*. (1)9: 174-178, pl. 19.
- HERRMANN, N. D., 2000. Dental analysis and palaeoecological assessment of *Baringa nelsonensis* (Marsupialia: Macropodidae: Macropodinae), and intermediate browsing/ grazing kangaroo from the Early Pleistocene Nelson Bay Formation, Victoria, Australia. *Unpublished Masters Thesis, Department of Earth Science, Monash University, Clayton, Victoria, Australia*.
- HOPE, J. H., 1978. Pleistocene mammal extinctions: The problem of Mungo and Menindee, New South Wales. *Alcheringa*. 2: 65-82.
- HOPE, J. H., LAMPERT, R. J., EDMONSON, E., SMITH, M. J. & VAN TETS, G. F., 1977. Late Pleistocene faunal remains from the Seton Rock Shelter, Kangaroo Island, South Australia. *Journal of Biogeography* 4: 363-385.
- HORTON, D. R., 1977. A 10,000 year old *Sarcophilus* from Cape York. *Search*. 10: 374-375.
- ILLIGER, J. K. W., 1811. C. Illigeri ...Prodromus systematis mammalium et avium additis terminis zoographicis utriusque classis, eorumque versione Germanica. *Berolini*.
- KENLEY, P. R. 1988. Southwestern Victoria. In *Geology of Victoria*, J. G. Douglas & J. A. Ferguson eds., Victorian Division, Geological Society of Australia Incorporated, Melbourne, 383-387.
- KERR, R., 1792. *The Animal Kingdom, or zoological system, of the celebrated Sir Charles Linnaeus; Class I. Mammalia, containing a complete systematic description, arrangement, and nomenclature, of all the known species and varieties of the Mammalia, or animals which give suck to their young: being a translation of that part of the Systema Naturae, as lately published, with great improvements, by Professor Gmelin of Goettingen*. J. Murray & R. Faulder, London, 1 Pt 1 xl 400 pp. 9 pls.
- LONG, J., ARCHER, M., FLANNERY, T. & HAND, S., 2002. *Prehistoric Mammals of Australia and New Guinea: One Hundred Million Years of Evolution*. University of New South Wales Press Ltd, University of New South Wales, Sydney.
- MACFADDEN, M. J., MACFADDEN, P., RICH, T. H., & WHITELAW, M. J., 1987. Magnetic Polarity Stratigraphy of the Pleistocene Section at Portland (Victoria) Australia. *Quaternary Research* 28: 364-373.
- MARSHALL, L. G., 1974. Late Pleistocene Mammals from the Keilor cranium site, southern Victoria, Australia. *Memoirs of the National Museum of Victoria* 35: 63-86.
- MARSHALL, L. G., 1973. Fossil vertebrate faunas from the Lake Victoria region of southwestern New South Wales. *Memoirs of the National Museum of Victoria* 34: 151-171.
- MERIGIAN, D., MURRAY, P. F., LAITZ, P. K. & JOHNSON, K. A., 2002. The Mygoora Local Fauna: a late Quaternary vertebrate assemblage from central Australia. *The Beagle: Records from the Museums and Art Galleries of the Northern Territory* 18: 77-93.
- MURRAY, P., 1991. The Pleistocene Megafauna of Australia. In *Vertebrate Palaeontology of Australasia*, P. Vickers-Rich, J. M. Monaghan, R. F. Baird, & T. H. Rich, eds. Monash Univer-

- sity Publishing Committee, Melbourne, 893-1070.
- OWEN, R. 1838. In *Three expeditions into the interior of eastern Australia, with descriptions of the recently explored region of Australia Felix, and of the present colony of New South Wales*, T. L. Mitchell, T & W Boone, London.
- OWEN, R., 1877. *Researches on the fossil remains of the extinct mammals of Australia; with a notice on the extinct marsupials of England, Volumes 1 & 2*. J. Erxleben, London.
- RICH, T. H., ARCHER, M., HAND, S. J., GODTHIELP, H., MUIRHEAD, J., PLEDGE, N. S., FLANNERY, T. F., WOODBURN, M. O., CASE, J. A., TEDFORD, R. H., TURNBULL, W. D., LUNDELIUS, E. L. *fil.*, RICH, L. S. V., WHITEHEAD, M. J., KEMP, A., & RICH, P. V., 1991. Appendix 1: Australian Mesozoic and Tertiary Terrestrial Mammal Localities. In *Vertebrate Palaeontology of Australasia*. P. Vickers-Rich, J. M. Monaghan, R. F. Baird, & T. H. Rich, eds. Monash University Publishing Committee, Melbourne. 1005-1070.
- RIDE, W. D. L., 1964. A review of the Australian fossil marsupials. *Journal of the Proceedings of the Royal Society of Western Australia*. 47: 97-131.
- SMITHSONIAN MUSEUM OF NATURAL HISTORY, DEPARTMENT OF SYSTEMATIC BIOLOGY, 1993. *Mammal Species of the World Database*. Smithsonian Institution, Washington D. C., United States of America. 1993. <http://nmnhwww.si.edu/msw/>
- SELWYN, A. R. C. 1858. On the geology of the gold-fields of Victoria. *Quarterly Journal of the Geological Society of London* 14: 533-538.
- SELWYN, A. R. C., 1860. Notes on the Geology of Victoria. *Quarterly Journal of the Geological Society of London* 16: 145-150.
- SRINIVASAN, M. S. & KENNET, J. P., 1981. A review of the Neogene planktonic foraminiferal biostratigraphy: Applications in the equatorial and South Pacific. *Society for Economic Palaeontologists and Mineralogists Special Publication* 32: 395-432.
- STIRTON, R. A., 1957. Tertiary Marsupials from Victoria, Australia. *Memoirs of the National Museum of Victoria* 21: 121-134.
- STRAHAN, R. (Ed.) 1998. *The Mammals of Australia*. Reed New Holland Publishers, Sydney.
- SZALAY, F. S., 1982. A new appraisal of marsupial phylogeny and classification. In *Carnivorous Marsupials Volume 2*. M. Archer ed., Surrey Beatty & Sons Pty. Ltd. and The Royal Zoological Society of New South Wales, Mosman, New South Wales, 621-640.
- THOMAS, O., 1903. Note on the technical name of the Tasmanian Devil. *Annals and Magazine of Natural History, series 7*, 11: 289.
- WELLS, R. T., 1978. Fossil mammals in the reconstruction of Quaternary environments with examples from the Australian fauna. In *Biology and Quaternary environments*, D. Walker & J. C. Guppy eds. Australian Academy of Science, Canberra, 103-204.
- WERDERLIN, L., 1987. Some observations on *Sarcophilus laniarius* and the evolution of *Sarcophilus*. *Records of the Queen Victoria Museum, Launceston* 90: 1-27.
- WHITE, M. A., 2002. Formation of the vertebrate bone beds of the Nelson Bay Formation, Portland, Victoria. *Unpublished Honours Thesis, Deakin University, Victoria*.
- WHITEHEAD, M. J., 1991. Magnetic polarity stratigraphy of Pliocene and Pleistocene fossil vertebrate localities in southeastern Australia. *Geological Society of America Bulletin* 103: 1493-1503.
- WROE, S., EBACH, M., AIHYONG, S., DE MUIZON, C., & MUIRHEAD, J., 2000. Cladistic analysis of Dasyuromorphian (Marsupialia) phylogeny using cranial and dental characters. *Journal of Mammalogy* 81(4): 1008-1024.

Manuscript received 16 June 2003

Revision accepted 17 November 2003



# SCOLOPENDRID CENTIPEDES CAUGHT BY PITFALL TRAPPING IN THE ULURU-KATA TJUTA NATIONAL PARK, NORTHERN TERRITORY, INCLUDING A RECORD OF AN APPARENT 'OUTBREAK'

ALAN L. YEN<sup>1,2</sup> & JOHN WAINER<sup>1,3</sup>

<sup>1</sup>Museum Victoria, GPO Box 666E, Melbourne, Victoria 3001, Australia

<sup>2</sup>Present address: Arthur Rylah Institute for Environmental Research,  
Department of Sustainability and Environment, PO Box 137, Heidelberg, Victoria  
3084, Australia

<sup>3</sup>Present address: Institute for Horticultural Development, Department of Primary  
Industry, Knoxfield, Private Bag 15, Ferntree Gully Delivery Centre, Victoria, 3156,  
Australia.

YEN, ALAN L. & WAINER, JOHN 2003:11:30. Scolopendrid centipedes caught by pitfall trapping in the Uluru-Kata Tjuta National Park, Northern Territory, including a record of an apparent 'outbreak'. *Proceedings of the Royal Society of Victoria* 115(2): 55-65. ISSN 0035-9211.

Pitfall trap surveys of fauna were conducted in the Uluru-Kata Tjuta National Park in 1995, 1997 and 2000. Relatively low numbers of scolopendrid centipedes were trapped during most of the surveys, but heavy summer rainfall in early 2000 may have been responsible for large numbers of individuals trapped then. The surveys have recorded a total of six species of scolopendrid centipedes from the Park, bringing the total known species there to seven.

*Key words:* Scolopendrid centipedes, Uluru-Kata Tjuta National Park

AUSTRALIAN centipedes are classified into five orders: the Scolopendrida, Geophilida, Lithobiida, Craterostigmatida and Scutigera (Harvey & Yen 1989). Scolopendrid centipedes are common in the drier regions of Australia. Fortyfour named species of scolopendrid centipedes are known from Australia, and they are predominantly solitary nocturnal predators (Edgecombe 2001) feeding on other invertebrates or occasionally on small vertebrates.

Recent faunal studies of both vertebrates and invertebrates in the Uluru-Kata Tjuta National Park have, under the joint management arrangement at the Park, involved participation by traditional owners (Anangu).

On the morning of 22 February 2000, drift fence pitfall traps that were running as part of a spinifex fire age faunal survey in the Uluru-Kata Tjuta National Park caught enormous numbers of scolopendrid centipedes – a total of 945 individuals in 54 traps (an average of 17.5 centipedes/trap)! The Uluru fauna surveys, conducted in 1995, 1997 and in October 2000, using the same trapping techniques but at different locations within the Park, found relatively small numbers of centipedes. This apparent 'outbreak' in numbers of centipedes was common

knowledge amongst the Anangu, who recognise several different colour forms, and had also been reported at the nearby tourist resort at Yulara. However, it had not been recorded in the scientific literature, and this was an opportunity to determine which species of scolopendrid centipedes occurred in the area, and whether particular species had these apparent spectacular increases in numbers.

In this paper we enumerate the scolopendrid centipedes trapped during the spinifex fire age faunal survey in February-March 2000 (SFS) in relation to sites and dates, and will compare results with the three Uluru fauna surveys (UFS) in 1995, 1997 and October 2000.

## METHODS

The Uluru fauna surveys were conducted in March-April 1995, September-October 1997 and October 2000. The fauna surveys involved eight sites (Fig. 1), and the same eight sites were sampled in 1995, 1997 and 2000 (Yen *et al.* 1996): UFS 1 (Open grassland to woodland at the base of Uluru), UFS 4 (Mulga shrubland), UFS 5 (Soft spinifex), UFS 6 (Hard spinifex), UFS 7 (Mallee shrubland), UFS 8 (Sandhill mulga), UFS14 (Hard spinifex transitional sandplain), and UFS15 (Alluvial fans and watercourse, Shrubland

to River Red Gum creekbed adjacent to Kata-Tjuta). The 2000 spinifex fire-age survey involved six sites (sites 1-6), each of which was divided into three classes of spinifex (older spinifex, younger spinifex, and ecotone). There were two replicates of each spinifex class at each site, resulting in six trap lines within each site. The term "spinifex" is used in this report to describe the various species of hummock grasses found in the study areas. The older spinifex sites (labeled A and B) were last burnt in 1976 or earlier. The younger spinifex sites (E and F) were burnt more recently. In all cases, there was an ecotone between the older and younger spinifex (labeled C and D). The SFS sites are shown in Fig. 1, and the dates on which the pitfall traps were run in 2000 were: SFS 5 (17-20 Feb), SFS 3 (21-24 Feb), SFS 2 (24-27 Feb), SFS 1 (27 Feb - 01 Mar), SFS 4 (01-04 Mar) and SFS 6 (04-07 Mar).

Centipedes were collected by drift fence pitfall trapping. At each site, the pitfall traps were arranged along two intersecting 25 metre lines arranged at right angles to each other, thus forming a cross. The traps were 20 litre buckets. One bucket was located at the intersection of the lines and four additional buckets along each line at six metre intervals. Consequently each pitfall trap site consisted of a set of nine buckets. A nylon drift fence was run along each line to connect the buckets. Within each sampling site, several replicate pitfall lines were operated concurrently. In both the UFSs and the SFS, the same pitfall trap line configuration was used; the only difference was that in the UFS, there were 4 sets of pitfall traps at each site, while in the SFS, there were 6 sets of pitfall traps. The traps were opened for three nights at each site, and examined in the morning and the evening. All invertebrates in the pitfall traps were collected with forceps and preserved in 70% ethanol. All pitfall trap material was eventually identified to the morphospecies level and lodged in the Museum of Victoria.

No attempt will be made in this paper to determine what environmental factors may be important in determining scolopendrid distribution in the UFS and the SFS sites. In the UFS, scolopendrid numbers, both in terms of species and individuals, are in the main low. Superficially, there did not seem to be a consistent trend of scolopendrid association with any of the major plant communities in the UFS. In the SFS, replicates were taken in older and younger spinifex, and ecotones, and the numbers of scolopendrids in these different aged spinifex is considered.

## RESULTS

The predominant family of centipedes collected in the pitfall traps was the Scolopendridae. A very small number of centipedes belonging to the Scutigerae were collected, but they have not been included in this paper. This dominance of the centipede fauna by the Scolopendridae was also found in the Camarvon Basin of Western Australia (Harvey *et al.* 2000). Unless otherwise stated, results from the UFS and the SFS are presented as number of individuals collected in each site, combining data from each of the pitfall lines. Consequently data from the UFS are based on 4 pitfall lines, while data from the SFS are based on 6 pitfall lines.

### *Fauna surveys 1995, 1997 and 2000*

The numbers of scolopendrids trapped was low in all three surveys (Table 1). Except for two occasions, fewer than 10 individuals were caught in any one pitline in any year. The exceptions were in 1995, when 24 *Ethmostigmus curtipes* were caught in pitline 3 at Site 15 and 20 *E. curtipes* in pitline 4 at Site 15. There was light rain on the night before these were trapped. While these two pitlines were adjacent, they were very different in vegetation: P3 was located in Ti-tree on sand and P4 on River Red Gum creek bed (primarily sand and rock).

In 1995, four species of scolopendrids were trapped, comprising a total of 121 individuals. They were dominated by two species: *Ethmostigmus curtipes* (67.8%) and *Scolopendra morsitans* (27.3%) (Table 1). In 1997, five species (84 individuals) were trapped. *E. curtipes* was again the dominant species, but much reduced in proportion (32.1%); *Scolopendra morsitans* (31.0%) was similar to 1995, while *Cormocephalus* sp. 2 (15.5%), *Cormocephalus* sp. 1 and *E. pachysoma* (10.7% each) occurred in higher proportions (Table 1). In the 2000 UFS, six species (69 individuals) were collected. The percentage compositions were similar to 1997 except that *E. pachysoma* was much reduced; *E. curtipes* (33.3%), *Scolopendra morsitans* (31.9%), *Cormocephalus* sp. 2 (15.9%) and *Cormocephalus* sp. 1 (14.5%) (Table 1).

The occurrence of each of the scolopendrid species thus varied in abundance and distribution over time and space. *Scolopendra morsitans* comprised approximately 33% of the individuals collected in each of the three faunal surveys; it occurred at all eight sites every survey with the exception of UFS 7 in 2000.






Fig. 1. Map of Uluru-Kata Tjuta National Park showing location of sites

*Cormocephalus* sp. 1 and 2 were either absent or very low in numbers in 1995, and comprised 10-15% of individuals collected in 1997 and 2000; over the 3 surveys, both species occurred in a total of five out of the eight UFS sites. *E. curtipes* ranged from 33-66% of the individuals trapped, and over the three surveys, was found in all eight UFS sites. In contrast, *E. pachysoma* and *E. rubripes* occurred in low numbers throughout the surveys and were only trapped in 3 of the UFS sites (Table 1).

The number of species trapped at any one site varied from one to five depending upon the year (Table 1). It is worth noting that over the three surveys, all six species were trapped in two sites (UFS7 and UFS8), five species at UFS5, four at UFS15, while only two species were trapped at UFS2 and three species at UFS1, UFS6 and UFS14 (Table 1).

#### *Spinifex* fire survey 2000

A total of 6 scolopendrid species, comprising 1859 individuals, was collected (Table 2). The fauna, based on percentage composition, was dominated by *E. pachysoma* (53.7%), *E. curtipes* (33.8%) and *Scolopendra morsitans* (11%). These three species were trapped at all six SFS sites, although their abundances, except for *Scolopendra morsitans*, varied between sites. *E. rubripes* was trapped in four of the SFS sites, while *Cormocephalus* sp. 1 and sp. 2 were trapped in three and two sites respectively (Table 2). In terms of relative abundances, *E. curtipes* dominated at four of the six SFS sites (SFS 1, 4, 5 and 6), *E. pachysoma* dominated at SFS3 and *Scolopendra morsitans* dominated at SFS5.

The sudden enormous increase in scolopendrid numbers caught in pitfall traps occurred at SFS3. Pitfall traps were opened at this site on the morning of 21 Feb and closed on the morning of 24 Feb. There was heavy rain on 19 Feb (36.5 mm) and 20 Feb (66.2 mm) before the traps at SFS3 were run (traps were SFS5 were open during the heavy rain). The pitfall traps had large numbers of centipedes on the morning of 22 Feb, followed by fewer on the mornings of Feb 23 and 24. The pitfall trap samples on the morning of 22 February were kept separate from those of the mornings of 23-24 February (when there were fewer centipedes). The data from SFS3 are presented in Table 3. The catch on 22 February was primarily *Ethmostigmus pachysoma*, between 124-180 individuals at each of the six pitfall lines. Small numbers of *E. curtipes* and *Scolopendra morsitans* were trapped on 22 February. The number of *E. curtipes* caught in

the traps was larger on 23-24 February, although the number of *E. pachysoma* trapped was marginally higher in three trap lines on these dates. In general, the major occurrence was the enormous increase in *E. pachysoma* numbers on the night of 21 February (and possibly very early in the morning of 22 February), and a sudden drop, although not a disappearance, on the nights of 22 and 23 February.

The SFS was conducted over 20 consecutive days. Considerable rainfall occurred on several days preceding the survey and also during the survey, and it is possible that this rain influenced centipede activity. The abundances of each species of centipede at each of the sites during the SFS are plotted in Fig. 2. *Scolopendra morsitans* was relatively steady in numbers at all sites during the SFS. *Cormocephalus* sp. 1 and sp. 2 occurred in low numbers after Feb 24. The responses of the three *Ethmostigmus* species differed; *E. curtipes* occurred in relatively high numbers throughout the survey, *E. pachysoma* occurred in very high numbers earlier during the survey, while *E. rubripes* occurred in relatively low numbers in four of the six sampling occasions.

The scolopendrid distribution between the different aged spinifex sites is presented in Table 4. More scolopendrids were trapped in older spinifex (A+B), followed by ecotones (C+D) and lowest numbers in younger spinifex (E+F). When examined on a species basis, the major differences in the different spinifex types is due to *Ethmostigmus curtipes*, which seems to prefer older spinifex over younger spinifex (Table 4).

## DISCUSSION

There are few previous records of scolopendrid centipedes from the Uluru-Kata Tjuta National Park. In his revisions of Australian scolopendrids, Koeh noted three species from within the Park: *Scolopendra laeta* Haase from Mt Olga and west gorges of Mt Olga (Koeh 1982), *Scolopendra morsitans* L. from Mt Olga (Koeh 1983a) and *Ethmostigmus rubripes* (Brandt) from Ayers Rock (Koeh 1983b). The UFS and the SFS added a further four species of scolopendrid, taking the number of recorded species to seven. Harvey *et al.* (2000) collected 13 species of scolopendrid centipedes by pitfall traps in the Camarvon Basin of Western Australia, although their study was conducted over a wider geographical region and they ran pitfall traps continuously over a 12 month period.






Fig. 2. Centipedes from spinifex fire survey

There are three issues to be discussed with regard to the scolopendrids trapped in the surveys at Uluru-Kata Tjuta. They are (1) the numbers of individuals collected by pitfall trap studies; (2) the response of the scolopendrids to heavy rain; and (3) the role of scolopendrids in the arid systems of Central Australia.

#### *Numbers of individuals and pitfall trapping*

In the few published studies on population densities of centipedes, the densities are generally low. The only scolopendrid species studied with regard to density are *Scolopendra amazonica* in Nigeria, which has a density of 0.16/m<sup>2</sup> (and a biomass of 140 mg/m<sup>2</sup>), although pitfall trapping was not the collecting method employed (Lewis 1972a). While it is difficult to estimate densities from pitfall trap studies, they can give an indication of whether numbers are low or high. In a fauna survey of the Mallee in north-western Victoria, drift fence pitfall trapping collected centipedes (Robertson *et al.* 1986), and a total of 1026 specimens, comprising 14 species, was collected (Yen & Lillywhite, unpublished data). However, the traps were spread over 122 sites, and each site comprised 10 x 20 litre pitfall traps which were run for five nights on five occasions over an 18 month period, giving an average of less than 2 individuals per pitfall line on each collecting occasion. Corey (1988) used pitfall traps in three plant communities over a six month period (detailed pitfall trap methodology were not given) in Florida (USA), and collected only 28 individuals from 5 species.

Throughout the UFS, the number of scolopendrid centipedes collected in pitfall traps was relatively low. The very large numbers were found at different sites during the SFS during a time of heavy summer rain. However, it appears that during the periods of no or minor rainfall, scolopendrid activity, and consequently catches in pitfall traps are low. The trap records from the Uluru fauna surveys suggest that the densities of scolopendrid centipedes are low.

#### *Response of centipedes to heavy rain*

However, the large numbers collected during the SFS suggest that scolopendrid densities may not be as low as suggested by results from the UFS. It is difficult to make definitive conclusions about this because (1) the SFS sites did not coincide with the UFS sites; and (2) it is not possible to distinguish site-specific dif-

ferences or the effects of heavy rain at the SFS sites because the pitfall traps at each site were run sequentially over time.

In Nigeria, the beginning of the rainy season initiated dispersal of young centipedes (Lewis 1972b). In South-East Asia, responses of two species of scolopendrid centipedes to water differed: *Scolopendra subspinipes* swims when immersed, while *Arrhabdotus octosulcatus* cannot swim and walks under water with its head out of the water. The latter is probably an arboreal species while the former is more likely to experience flooding from tropical down pours (Lewis 1980). Finally, flushing with water has been used as a collecting technique for centipedes (Lewis 1972a).

The results suggest that the increased activity of centipedes, and consequently the larger total numbers trapped, during the SFS is due to heavy rainfall. However, the responses seem to be related to the individual species, especially *Ethmostigmus pachysoma*. The numbers of *Scolopendra morsitans* and *Ethmostigmus curtipes* were steady throughout the SFS, while the numbers of both *Cormocephalus* species and *Ethmostigmus rubripes* remained relatively low. It is suggested that the generally higher rainfall may be responsible for the increased numbers of centipedes caught in the pitfall traps, but the rain may have affected some species more than others. The large numbers of *E. pachysoma* occurred on Feb 21-24 at site SFS3, when the numbers of *Scolopendra morsitans* was lowest and no *Cormocephalus* species or *Ethmostigmus rubripes* were trapped. This may indicate that the last three species did not occur at SFS3, that *E. pachysoma* and *E. curtipes* have higher densities at this site, and the recorded response was due to the heavy rain in the days preceding the trapping. There was even heavier rainfall on 11-13 February, before the SFS began, and it is possible that any response to heavy rain by centipedes may be rapid and short.

Anecdotal information from traditional owners and from visitors to the area indicate previous increased centipede activity after heavy rains. Whether the response is a cue for some aspect of centipede behaviour (dispersal, mate seeking, etc) is not known, although most individuals collected seemed to be mature. Alternatively, the increased activity may simply have been due to centipedes being flushed out of their subterranean habitat.



1995 Uluru fauna survey												
S1	S4	S5	S6	S7	S8	S14	S15	Total	%			
<i>Scolopendra morsitans</i>	4	2	11	2	1	7	4	2	33	27.3		
<i>Cormocephalus</i> sp. 1	0	0	4	0	0	0	0	0	4	3.3		
<i>Cormocephalus</i> sp. 2	0	0	0	0	0	0	0	0	0	0		
<i>Ethmostigmus curtipes</i>	6	5	1	0	0	4	0	66	82	67.8		
<i>Ethmostigmus pachysoma</i>	0	0	0	0	0	0	0	0	0	0		
<i>Ethmostigmus rubripes</i>	0	0	0	0	1	1	0	0	2	1.7		
Total number individuals	10	7	16	2	2	12	4	68	121			
Number species	2	2	3	1	2	3	1	2	4			

1997 Uluru fauna survey												
S1	S4	S5	S6	S7	S8	S14	S15	Total	%			
<i>Scolopendra morsitans</i>	2	2	6	5	2	2	4	3	26	31		
<i>Cormocephalus</i> sp. 1	3	0	4	0	0	0	0	2	9	10.7		
<i>Cormocephalus</i> sp. 2	0	0	0	7	1	0	5	0	13	15.5		
<i>Ethmostigmus curtipes</i>	4	0	8	3	0	7	5	0	27	32.1		
<i>Ethmostigmus pachysoma</i>	0	0	1	0	3	5	0	0	9	10.7		
<i>Ethmostigmus rubripes</i>	0	0	0	0	0	0	0	0	0	0		
Total number individuals	9	2	19	15	6	14	14	5	84			
Number species	3	1	4	3	3	3	3	2	5			

2000 Uluru fauna survey												
S1	S4	S5	S6	S7	S8	S14	S15	Total	%			
<i>Scolopendra morsitans</i>	3	3	5	2	0	3	3	22	31.9			
<i>Cormocephalus</i> sp. 1	1	0	2	0	4	1	0	2	10	14.5		
<i>Cormocephalus</i> sp. 2	0	0	3	0	1	2	5	0	11	15.9		
<i>Ethmostigmus curtipes</i>	4	0	0	1	3	4	4	7	23	33.3		
<i>Ethmostigmus pachysoma</i>	0	0	1	0	0	0	0	0	1	1.4		
<i>Ethmostigmus rubripes</i>	0	0	0	0	0	1	0	1	2	2.9		
Total number individuals	8	3	11	3	8	11	12	13	69			
Number species	3	1	4	2	3	5	3	4	6			

Table 1. Number of species and individuals of scolopendrid centipedes trapped in the 1995, 1997 and 2000 Uluru fauna surveys

*The role of scolopendrids in the arid systems of Central Australia*

If we accept that centipede activity is low during times of non-rainfall or light rainfall, and that heavy rainfall flushes out large numbers, then the densities of scolopendrids living in the sand may be much higher than we imagine. Lewis (1981) reported that centipedes can survive aridity by burrowing deep into the soil. Anangu observed centipede burrows in the sand at Uluru. The question arises as to whether they form structured burrows or whether they simply 'swim' in sand. Other interesting topics include how frequently they emerge above ground, and whether they catch food in their subterranean habitat. *Scolopendra* and *Cormocephalus* are solitary species (Lewis 1981).

It is interesting to speculate upon the ecological role that centipedes play in the semi-arid and arid zones of Australia. Intraspecific encounters are usually aggressive (Edgecombe 2001). They are relatively large predators (in invertebrate terms), and as such would be expected to play a significant role in view of the large numbers trapped during the SFS. A wide variation in the diets of scolopendrids has been recorded from around the world; food ranges from invertebrates to small vertebrates (Lewis 1981). There is an abundance of potential invertebrate food for centipedes in Central Australia. The ground-active invertebrate fauna is dominated by ants and termites, and Anangu report that during dry conditions, centipedes catch ants as food. After rain, they emerge to catch termites. In turn, their relatively large body size would provide a decent meal for higher level predators, and Anangu report that they are eaten by sand goannas.

In the western desert region of Australia, centipedes are called 'wanatjiti' or 'kanparka' in the Pitjantjatjara language. According to the traditional owners (Anangu), there is Tjukurpa (Law) associated with centipedes; and some of this information is not available.

Much remains unknown about scolopendrid centipedes. Their basic life histories and longevity remains to be determined. Population numbers and fluctuations and habitat factors that influence their distribution are still unknown.

#### ACKNOWLEDGEMENTS

We wish to thank Peter Lillywhite (Museum Victoria) and Greg Edgecombe (Australian Museum) for

	SFS1	SFS2	SFS3	SFS4	SFS5	SFS6	Total	%
<i>Scolopendra morsitans</i>	41	30	23	38	35	37	204	11
<i>Cormocephalus</i> sp. 1	0	1	0	8	0	5	14	0.8
<i>Cormocephalus</i> sp. 2	1	0	0	4	0	0	5	0.3
<i>Ethmostigmus curtipes</i>	183	24	131	181	64	46	629	33.8
<i>Ethmostigmus pachysoma</i>	6	1	965	2	11	14	999	53.7
<i>Ethmostigmus rubripes</i>	2	2	0	0	1	3	8	0.4
Total number individuals	233	58	1119	233	111	105	1859	
Number species	5	5	3	5	4	5	6	

Table 2. Number of species and individuals of scolopendrid centipedes trapped during spinifex fire age survey Feb-March 2000



assistance with identification of the centipedes, and Leigh Ahern for field assistance during the 2000 UFS. The work was conducted with financial support from Environment Australia, and we are indebted to Jake Gillen and Sam Rando for their organization of the project. Various staff members of Parks Australia North assisted with various aspects of these surveys, and traditional owners (Imantura, Daisey Walkabout, Joyce Tjalyiri, Ginger Tjintalka and Billywara) also provided support and information.

Julian Reid and Steve McAlpin, the vertebrate zoologists on these surveys, also provided assistance. The work was conducted under the conditions of Environment Australia Scientific Research Permit No. RU-2000-4.

# REFERENCES

- COREY, D.T. 1988. Centipedes and millipedes in three Central Florida plant communities (Chilopoda and Diplopoda). *Florida Entomologist* 71:659-662.
- EDGEcombe, G. 2001. Centipedes: the great Australian bite. *Nature Australia* 26:42-51.
- HARVEY, M.S., SAMPEY, A., WEST, P.L.J. & WALDOCK, J.M. 2000. The Chilopoda and Diplopoda of the southern Carnarvon Basin, Western Australia. *Records of the Western Australian Museum Supplement* No. 61:323-333.
- HARVEY, M.S. & YEN, A.L. 1989. *Worms to Wasps: An Illustrated Guide to Australia's Terrestrial Invertebrates*. Oxford University Press: Melbourne. 203 pp.
- KOCH, L.E. 1982. Taxonomy of the centipede *Scolopendra laeta* Haase (Chilopoda: Scolopendridae). *Zoological Journal of the Linnean Society* 76:125-140.
- KOCH, L.E. 1983a. Morphological characters of Australian scolopendrid centipedes, and the taxonomy and distribution of *Scolopendra morsitans* L. (Chilopoda: Scolopendridae: Scolopendrinae). *Australian Journal of Zoology* 31:79-91.
- KOCH, L.E. 1983b. A taxonomic study of the centipede genus *Ethmostigmus* Pocock (Chilopoda: Scolopendridae: Otostigminae) in Australia. *Australian Journal of Zoology* 31:835-849.
- LEWIS, J.G.E. 1972a. The population density and biomass of the centipede *Scolopendra amazonica* (Bücherl) (Scolopendromorpha: Scolopendridae) in Sahel. *Entomologist's Monthly Magazine* 108:16-18.

	Site 3A		Site 3B		Site 3C		Site 3D		Site 3E		Site 3F	
	22-Feb	23/24 Feb	22-Feb	23/24 Feb	22-Feb	23/24 Feb	22-Feb	23/24 Feb	22-Feb	23/24 Feb	22-Feb	23/24 Feb
<i>Scolopendra morsitans</i>	1	0	2	5	3	1	2	1	1	1	3	3
<i>Ethmostigmus curtipes</i>	8	36	16	31	4	15	1	0	4	8	1	7
<i>Ethmostigmus pachysoma</i>	166	22	180	5	124	7	144	11	151	10	134	10

Table 3. Pitfall trap catches of scolopendrids at site 3 during 'outbreak' in February 2000. he traps were opened on 21 Feb and closed on 24 Feb.

SFS Site and date	Species	Old	Ecotone	Young
Site 1. 27 Feb - 01 Mar 2000	<i>Scolopendra morsitans</i>	13	20	8
	<i>Cormocephalus</i> sp. 1	0	0	0
	<i>Cormocephalus</i> sp. 2	1	0	0
	<i>Ethmostigmus curtipes</i>	92	72	19
	<i>Ethmostigmus pachysoma</i>	3	2	1
	<i>Ethmostigmus rubripes</i>	0	1	1
	Total	109	95	29
Site 2. 24-27 Feb 2000	<i>Scolopendra morsitans</i>	4	14	12
	<i>Cormocephalus</i> sp. 1	0	0	1
	<i>Cormocephalus</i> sp. 2	0	0	0
	<i>Ethmostigmus curtipes</i>	9	8	7
	<i>Ethmostigmus pachysoma</i>	0	0	1
	<i>Ethmostigmus rubripes</i>	0	1	1
	Total	13	23	22
Site 3. 21-24 Feb 2000	<i>Scolopendra morsitans</i>	8	7	8
	<i>Cormocephalus</i> sp. 1	0	0	0
	<i>Cormocephalus</i> sp. 2	0	0	0
	<i>Ethmostigmus curtipes</i>	91	20	20
	<i>Ethmostigmus pachysoma</i>	373	287	305
	<i>Ethmostigmus rubripes</i>	0	0	0
	Total	472	314	333
Site 4. 01-04 Mar 2000	<i>Scolopendra morsitans</i>	9	15	14
	<i>Cormocephalus</i> sp. 1	5	1	2
	<i>Cormocephalus</i> sp. 2	1	0	3
	<i>Ethmostigmus curtipes</i>	106	33	42
	<i>Ethmostigmus pachysoma</i>	1	0	1
	<i>Ethmostigmus rubripes</i>	0	0	0
	Total	122	49	62
Site 5. 17-20 Feb 2000	<i>Scolopendra morsitans</i>	16	10	9
	<i>Cormocephalus</i> sp. 1	0	0	0
	<i>Cormocephalus</i> sp. 2	0	0	0
	<i>Ethmostigmus curtipes</i>	13	45	6
	<i>Ethmostigmus pachysoma</i>	6	1	4
	<i>Ethmostigmus rubripes</i>	0	0	1
	Total	35	56	20
Site 6. 04-07 Mar 2000	<i>Scolopendra morsitans</i>	18	15	4
	<i>Cormocephalus</i> sp. 1	4	0	1
	<i>Cormocephalus</i> sp. 2	0	0	0
	<i>Ethmostigmus curtipes</i>	21	24	1
	<i>Ethmostigmus pachysoma</i>	2	9	3
	<i>Ethmostigmus rubripes</i>	0	1	2
	Total	45	49	11
	Total	45	49	11
Total for all sites	<i>Scolopendra morsitans</i>	68	81	55
	<i>Cormocephalus</i> sp. 1	9	1	4
	<i>Cormocephalus</i> sp. 2	2	0	4
	<i>Ethmostigmus curtipes</i>	332	202	95
	<i>Ethmostigmus pachysoma</i>	385	299	315
	<i>Ethmostigmus rubripes</i>	0	3	5
	Total	796	586	478

Table 4. Differences between old, ecotone and young spinifex sites



- LEWIS, J.G.E. 1972b. The life histories and distribution of the centipedes *Rhysida munda togoensis* and *Ethmostigmus trigonopodus* (Scolopendromorpha: Scolopendridae) in Nigeria. *Journal of Zoology* 167:399-414.
- LEWIS, J.G.E. 1980. Swimming in the centipede *Scolopendra subspinipes* Leach (Chilopoda, Scolopendromorpha). *Entomologist's Monthly Magazine* 116:121-122.
- LEWIS, J.G.E. 1981. *The Biology of Centipedes*. Cambridge University Press: Cambridge. 476 pp.
- ROBERTSON, P., BENNETT, A.F., LUMSDEN, L.F., SILVEIRA, C.E., JOHNSON, P.G., YEN, A.L., MILLEDGE, G.A., LILLYWHITE, P.K. & PRIBBLE, H.J. 1989. Fauna of the Mallee study area, north-western Victoria. *Arthur Rylah Institute for Environmental Research Technical Report Series No.* 87. 91 pp.
- YEN, A.L., REID, J.R.W. & WAINER, J. 1996. A pitfall trap survey of ground dwelling invertebrates. In *Monitoring the Vertebrate Fauna of Uluru National Park, Phase II - Final Report*. Reid, J.R.W. & Hobbs, T.J. eds., Consultancy report to the Australian Nature Conservation Agency. CSIRO, Alice Springs. pp. 139-159.

Manuscript received 01 May 2003

Revision accepted 19 October 2003





# THE BALLARAT METEORITE, A FOSSIL IAB IRON FROM VICTORIA, AUSTRALIA

WILLIAM D. BIRCH<sup>1</sup> & LEONARD E. SAMUELS<sup>2</sup>

<sup>1</sup>Museum Victoria, GPO Box 666E, Melbourne, Victoria, 3001, Australia

<sup>2</sup>Department of Materials Engineering, Monash University, Clayton, Victoria, 3168, Australia

BIRCH, W. D. & SAMUELS, L. E., 2003:11:30, The Ballarat meteorite, a fossil IAB iron from Victoria, Australia. *Proceedings of the Royal Society of Victoria* 115(2): 67-75. ISSN 0035-9211.

The Ballarat meteorite was discovered in a palaeoplacer ('deep lead') deposit during gold mining in the Park Company leases at Ballarat, Victoria, during the late 1860s to early 1870s. The meteorite is a 12–15 g fragment from a type IAB iron that underwent disaggregation due to weathering and transport in a vigorous stream environment. The meteorite consists essentially of kamacite with a subgrain texture marked by partially resorbed Neuman bands and with plentiful schreibersite inclusions. The average nickel content is in the range 6.1–6.3 wt. % as determined by EPMA and INAA methods and with high Ge and Ga contents for the type. The maximum age of the meteorite is unconstrained but it is at least 3 million years old, as this is the age of the basalt flows that buried the palaeoplacer deposit immediately west of Ballarat. Reconstruction of the prebasaltic landscape indicates that the meteorite fell close to the drainage divide at the time and has probably not travelled far. The specimen has been preserved due to the reducing conditions maintained in the deep lead environment.

*Key words:* Ballarat, fossil meteorite, palaeoplacer, Victoria, IAB iron

THE BALLARAT meteorite is a new nickel-poor IAB iron found during 19<sup>th</sup> Century mining operations at Ballarat, Victoria, Australia. A single fragment, likely to have been about 15 g originally, was discovered in a buried palaeoplacer deposit in the Park Company gold mine between 1867 and 1874. The circumstances of its discovery are not known, but it appears to have found its way into the mineral collection of Henry Rosales (1820–1916), a Spanish-born, German-trained mining engineer, who was engaged in contract work in the region at the time. Rosales donated his mineral collection to the University of Melbourne in the 1890s. Many of his samples have original hand-written labels but, unfortunately, that for the meteorite fragment is missing. While its provenance may therefore be questioned, the overall documentation of the Rosales collection and the historical connections between Ballarat, Rosales and documented discoveries in the Park Company mine suggest the meteorite is authentic. The historical mineral collections of the University were transferred to Museum Victoria in the late 1980s and the meteorite was rediscovered some years later. At an unknown time, a small portion of the sample

has been removed for analysis, which detected nickel. Whether this was done by Rosales at the Ballarat School of Mines laboratory is not known, but the sample was registered in the University collection as 'meteoric (sic) iron'. Following further investigation by the present authors, the Ballarat meteorite was approved by the Nomenclature Committee of the Meteoritical Society. It is catalogued as specimen E15649 in Museum Victoria's meteorite collection.

## *The discovery site*

Rich shallow alluvial gold deposits were discovered in the Ballarat region in central Victoria in 1851. From the mid-1850s, mining began to follow the deposits deeper beneath younger sediments in the eastern part of the field and basalt flows to the west (Baragwanath 1923; Canavan 1988). Two major deep lead systems, the Inkerman and Golden Point leads, are buried by up to four lava flows of the Newer Volcanics, in places over 100 m thick in total (Fig. 1). Various companies took out leases and sank shafts to exploit the alluvial deposits. The Park Company was formed in November 1867 and by using several former companies'

shafts, opened up rich alluvial deposits in so-called "reef wash", at depths between about 250 and 320 feet (75–100 m). These were in an area between the Inkerman and Golden Point leads to the north and southeast, respectively, and about 30 metres above them (Baragwanath 1946). Until it closed in March 1874, the mine produced 95 000 ounces of gold.

Although the precise location within the Park Company mine is not known, the coordinates for the discovery site of the Ballarat meteorite would be close to 37° 34' 15" S; 143° 49' 55" E.

Several other metallic minerals were collected from the Park Company lead at about the same time. Rolled fragments of native copper and galena were recorded by Krausé (1882, 1896). Galena and pyrite specimens from the mine were exhibited in the Philadelphia Exhibition of 1875/6. While there is no specific mention of 'native iron' being found, it appears that some systematic collecting of unusual metallic minerals was undertaken while the mine was operating.

### *Features and composition of the meteorite*

The surviving piece of the Ballarat meteorite is a rough flattened mass of about 10 g, with dimensions 20 x 15 x 5 mm (Fig. 2). A fragment of about 1.5 g was removed to prepare a polished section and one of about 0.25 g for instrumental neutron activation analysis (INAA). There is only a very thin rust coating on the bright metal, which shows no propensity to alter under ambient conditions. The hackly surface of the piece suggests it is a fragment broken from a larger crystalline mass of kamacite, probably during transport.

In polished section the texture shows subgrains of kamacite crossed by partly resorbed Neumann bands and with scattered roughly prismatic crystals of

schreibersite ('rhabdite') up to about 0.05 mm long (Fig. 3). In detail these have ragged margins marked by an unknown precipitate (possibly earlsbergite or, less likely, roaldite) that appears to represent an incipient reaction halo with the enclosing kamacite (Fig. 4). These features closely resemble those of the rhabdites described from the Veevers IIAB iron by Bevan *et al.* (1995). Throughout the matrix kamacite and along the boundaries of Neumann bands are minute grains of another unknown phase. One outer portion of the fragment shows markedly deformed features, suggestive of a strong localised impact during transport. Rhabdite crystals have been bent along shear planes in the deformed section. The boundary between this deformed region and the main kamacite body is possibly a fragmented schreibersite grain (Fig. 5).

The meteorite was analysed for major elements (Fe, Ni, Co, P) using an electron microprobe and for Ni and minor elements by INAA. The average Ni content from 22 microprobe analyses is 6.1 wt % (range 5.6–6.6), slightly lower than the 6.3 wt % obtained by the INAA method (see Table 1). Ga, Ge and Ir contents of 95, 480 and 1.1 i g/g, respectively, indicate that Ballarat is a low Ni, high-Ga and high-Ge IAB iron, rather than a IIAB iron.

Analyses of seven schreibersite grains gave a formula of  $\text{Fe}_{1.64}\text{Ni}_{1.36}\text{P}$ .

### GEOLOGICAL CONTROLS ON AGE AND PRESERVATION

The palaeoplacer in which the Ballarat meteorite was found forms part of an extensive Cainozoic fluvial system in the West Victorian Uplands, in which rivers flowed north and south from an E–W-trending divide (Canavan 1988). The oldest palaeoplacers are

	Fe wt%	Ni wt%	Co wt%	P wt%	Ga ?g/g	Ge ?g/g	Ir ?g/g
Microprobe	94.17	6.10	0.36	0.035			
INAA		6.3			95	480	1.1

Table 1: Chemical analyses of the Ballarat meteorite





Fig. 1. Simplified geological map of the Ballarat district (after Finlay & Douglas, 1992). Main deep leads are shown as dotted lines. Lava flows from Mt Rowan eruption point cover the deep leads to the west of Ballarat, including the leases operated by the Park Company (see Fig. 6 for detail indicated by rectangle).





Fig. 2. The fragment of the Ballarat meteorite remaining after samples were removed for microscopy and analysis. The piece is 20 mm long. Museum Victoria specimen E15649.

unfossiliferous sheets of coarse quartz gravels known as the White Hills Gravel. These were laid down, probably in the very early Cainozoic, in broad shallow valleys cut into a landscape that had undergone extensive weathering. Later, younger, narrower valleys began to develop within the older broad valleys containing the White Hills Gravel. As this system advanced, streams eroded through the blanketing White Hills Gravel and incised valleys into the Palaeozoic bedrock, leaving remnant patches of gravel perched on interfluvies. The conglomerates, sands, clays and lignites laid down in these new valleys were to become known as 'deep leads', which in many parts of the region were extremely rich in secondary gold. These palaeoplacers were deposited at different times and at different elevations, during cycles of down-cutting and back-filling. Many were subsequently buried by Pliocene and younger basaltic lava flows. Lack of fossils, later faulting and inadequate outcrop mean that the palaeoplacers are difficult to date and

correlate. Dating of rare pollen remains shows that some of these deep leads began to develop as early as the late Eocene. However, deposition was long-lived and continued right through until the Pliocene, when extensive volcanism caused disruption of the palaeodrainage systems (Hughes & Carey 2002).

In the Ballarat goldfield, there are several types of gold-bearing palaeoplacers. Baragwanath (1923) considered that pre-basaltic 'deep lead reef wash', such as was worked in the Park Company leases in the western part of the field, was probably equivalent to remnant patches of gravel outcropping on low hills to the east. These exposed patches are now regarded as the early Cainozoic White Hills Gravel (Taylor et al. 1996). Historical descriptions of the buried reef wash describe a similar lithology of well-water-worn pebbles and boulders of quartz, some of enormous size, with subordinate fragments of bedrock (slate and sandstone) and pieces of water-worn siliceous cemented material from older deposits. Typically, reef wash



Fig. 3. General texture of the Ballarat meteorite showing subgrain boundaries in kamacite with partly resorbed Neumann bands and crystals of schreibersite. Field of view is about 1.5 mm across.

deposits were spread over an expanse of flat or undulating bedrock above the normal level of the main stream channel or 'gutter' (Hunter, 1909). Channels up to 60 m deep were cut through the reef wash by later streams and filled with 'gutter wash' which, after burial by lava flows, form the deep leads proper.

The palaeodrainage system in the vicinity of the meteorite's discovery has been investigated by Taylor & Gentle (2002), using bore-hole data and mine records. A reconstruction of the prebasaltic bedrock topography suggests that a 'deep lead divide' separated the Inkerman and Golden Point leads. Trending generally NE–SW, the divide passed through the area mined by Park Company (Fig. 6). It appears that the reef wash the company removed came mainly from two poorly delineated remnants of an older fluvial system preserved on either side, but close to the top, of the divide.


This geological environment provides some constraints on the timing and location of the fall of the Ballarat meteorite. Based on the reconstruction of Taylor & Gentle (2002), the divide on which the meteorite-bearing gravels were perched immediately before the basaltic eruptions that buried them appears to have been close to the main continental divide when the White Hills Gravel was deposited in the early Cainozoic. This restricts the fall site of the meteorite to somewhere on or near this palaeodivide and therefore not far from its final resting-place in the Park Company leases. This is the case regardless of whether the 'reef wash' gravels in which the meteorite was found represent a patch of the White Hills Gravel preserved beneath the basalt, or a younger deposit associated with the early stages of deep lead formation.



Fig. 4. Schreibersite prism showing incipient reaction halo with the enclosing kamacite. Field of view is about 0.08 mm across.

Such a distinction does, however, affect estimates of the maximum age of the fall. If the reef wash in the Park Company leases represents White Hills Gravel, then the meteorite could be as old or older than late Eocene, based on the time of deposition estimated for the White Hills Gravel by Hughes & Carey (2002). It could have fallen onto the Mesozoic surface that predates the White Hills Gravel before being incorporated into the deposits (Fig. 7). The minimum age of the fall is constrained by radiometric dates for basalts from the Ballarat region, which are between 2 and 4 Ma (Taylor et al. 1996; King 1985). The most representative age of 2.5–3 Ma is for lava from Mt Rowan, a volcano about 4 km to the northwest of the Park Company leases. This eruption point is the most likely

Fig. 5. Possible fragmented schreibersite grain along the interface between main body of fragment and deformed portion, marked by shear-bound etch markings. Field of view is 1.5 mm across.







Fig. 6. Reconstruction of the prebasaltic topography in the vicinity of the Park Company leases, modified from Taylor & Gentle (2002). The positions of the areas of gravel wash mined by the Park Company are based on the geological map of Baragwanath (1923).





*Fig. 7.* Possible simplified history of the Ballarat meteorite in terms of the late Mesozoic – Cainozoic landscape evolution in the Ballarat district (after Taylor & Gentle, 2002). Having fallen on the weathered palaeosurface, possibly as early as the late Mesozoic, the meteorite was corroded and broken up during short-distance transport and burial. Subsequently, deep lead valleys were incised through the higher level alluvial deposits. Lava flows (3 Ma) buried these gravels and filled the deep leads.

source of many of the flows that buried the deep leads, including the uppermost flows at Alfredtown, less than 2 km west of the Park Company leases (Fig. 1).

The original size of the meteorite cannot be estimated, but it is likely that the surviving piece is from a much larger mass. At times, the fluvial deposits would have been high-energy environments, as indicated by the abundance of pebbles of massive reef quartz. Under these 'ball-mill' conditions, a relatively coarse-grained iron meteorite would tend to respond to impacts by fracturing along grain or subgrain boundaries, probably weakened by incipient oxidation, with adjacent regions being deformed. Schreibersite grains oriented along subgrain boundaries may have facilitated this fracturing (Fig. 5). Features of the Ballarat meteorite are consistent with such treatment and response.

Once buried by the lava flows, the placer deposits would have been immobilised and conditions would gradually have become reducing, particularly if organic matter was present. The widespread presence of pyrite layers in deep lead gravels indicates they were generally reducing environments. In this situation, a piece of meteoritic iron enclosed in the deposits was likely to survive. The reported presence of other metallic fragments, such as unoxidised copper metal and galena, in the Park Company lead is consistent with reducing conditions.

## CONCLUSIONS

While the full circumstances surrounding the discovery and preservation of the Ballarat iron are unrecorded, there are sufficient links between the various strains of evidence to suggest it is a genuine occurrence of a fossil meteorite. It represents a kamacite plate or bundle of plates broken from a coarse-grained octahedrite. The microstructure of partially resorbed Neumann bands and incipient reaction haloes around rhabdite grains is evidence for mild annealing in a low-magnitude, short-lived heating episode.

Sometime before 3 Ma, the precursor mass of the Ballarat meteorite fell close to the main drainage divide in the Ballarat region. Following an unknown period of weathering, the meteorite was caught up in a high-energy fluvial environment that broke it into fragments. These were carried a short distance to be deposited in a placer deposit that was subsequently buried by 3-Ma basalt flows. At least one fragment survived in this environment to be collected during

mining operations between 1867 and 1874.

The Ballarat meteorite (IAB) is one of the most common types of iron meteorite, with over 120 known examples. Yet its small size and the circumstances of discovery make it highly unusual. It is the smallest iron meteorite known from Victoria, supplanting Wedderburn (30 g) found about 125 km to the north of Ballarat in 1951. Possibly the only smaller iron meteorite known is Castray River, from Tasmania, with a total mass of about 10 g (Bevan, personal communication).

## ACKNOWLEDGEMENTS

The authors wish to thank John Wasson (University of California, Los Angeles) for determining the composition of the Ballarat meteorite by INAA. The manuscript was kindly read by Martin Hughes (CSIRO Exploration and Mining) and David Taylor (Geological Survey of Victoria), who provided helpful advice on the age and origin of the palaeoplacer deposits in the Ballarat district, and by Alex Bevan (Western Australian Museum), who advised on the classification and phases present. David Friedman (Faculty of Engineering, RMIT University) assisted with cutting the meteorite.

## REFERENCES

- Baragwanath, W., 1946. Gold and Minerals. Geological Survey of Victoria (unpublished).
- Baragwanath, W., 1923. The Ballarat Goldfield. *Memiors of the Geological Survey of Victoria* 14.
- Bevan, A. W. R., Shoemaker, E. M. & Shoemaker, C. S., 1995. Metallography and thermo-mechanical treatment of the Veevers (IAB) crater-forming iron meteorite. *Records of the Western Australian Museum* 17: 51–59.
- Canavan, F., 1988. Deep lead gold deposits of Victoria. *Bulletins of the Geological Survey of Victoria* 62.
- Finlay, I. S. & Douglas, P. M., 1992. Ballarat mines and deep leads. 1:100,000 map. *Geological Survey of Victoria Report* No. 94.
- Hughes, M. J. & Carey, S. P., 2002. Palaeodrainage development of the West Victorian Uplands, Victoria, Australia. *Published papers, Victoria Undercover Conference, 2002*: 155–159.

- Hunter, S. B., 1909. The Deep Leads of Victoria. *Memoirs of the Geological Survey of Victoria* 7.
- King, R. L., 1985. Explanatory notes on the Ballarat 1:250 000 geological map. *Geological Survey of Victoria Report* No. 75.
- Krausé, F. M., 1882. Museum and Library Report. *Annual Report of the Ballarat School of Mines*.
- Krausé, F. M., 1896. An introduction to the study of mineralogy for Australian readers. Robertson and Company, Melbourne, 352 pp.
- Taylor, D.H., Whitehead, M. L., Olshina, A. & Leonard, J.G., 1996. Ballarat 1:100 000 map sheet geological report. *Geological Survey of Victoria Report* No. 101.
- Taylor D. H. & Gentle, L. V., 2002. Evolution of the deep-lead palaeodrainages and gold exploration at Ballarat, Australia. *Australian Journal of Earth Sciences* 49(5): 869–878.

Manuscript received 21 April 2003

Revision accepted 9 December 2003





TRANSACTION  
OF THE  
ROYAL SOCIETY OF VICTORIA





**LECTURE TO THE DANISH CLUB**  
**18<sup>th</sup> September, 2003**  
**by Phillip Law**

I accepted the invitation to lecture here tonight – a very rare acceptance – because I wanted to pay a tribute to the part played by Denmark in the development of Australia's role in Antarctic affairs.

At a time when Australia's plans to establish a permanent station in Antarctica were stalled through lack of a suitable ship, Denmark provided the answer to the problem in the form of a small icebreaker "Kista Dan". In December 1953, her departure from Melbourne with the pioneer Mawson party marked the beginning of a 34-year association with the Danish shipping company "J. Lauritzen" from which we chartered, in succession, four ships.

I was honoured to become a friend of the head of the company – Knud Lauritzen – who on several occasions entertained me and my wife Nel during our several visits to Copenhagen. One entertainment was particularly memorable. Our visit coincided with a visit to Copenhagen by the Royal

Covent Garden Ballet Company from London. Lauritzen had arranged a bay cruise for the ballet dancers on board the company's small training sailing barque "Lilla Dan" and had invited Nel and me to join them.

It was a perfect warm, sunny day. We sailed along, sunning ourselves on deck and partaking of oysters, prawns, crayfish and champagne. A piano accordion player provided jazz music and the ballet dancers jived enthusiastically. It was really magic!

*SOME ANTARCTIC LEADERS*

When one comes to look at what has been written about the history of Antarctic exploration, one is not very impressed. The trouble is that certain patterns of thought concerning Antarctic events become widely established as a result of the media's portrayal of them and the effect of such portrayal upon the general public. Long before some historian comes along to write up the events, they



have become enveloped in shrouds of ignorance, error, emotion, sentimentality, chauvinism and exaggeration.

Tonight I look back on the history of Antarctic exploration to compare the accomplishments and leadership qualities of some of the world's great Antarctic expeditioners and, in doing so, to show how their reputations confirm my opening statement.

Those I shall discuss are Scott, Shackleton, Amundsen and Mawson.

But first, I should like to stress the difference between discovery and exploration. Discovery, in the Antarctic context, is seeing geographical features for the first time. Exploration, in that context, is describing the features, and accurately determining their locations so they can be mapped. This is a most important distinction and one that, unfortunately, has seldom been made.

In assessing the accomplishments of these men, I shall use four criteria :

- i) what they, personally, explored and discovered;
- ii) what the men under their leadership achieved;
- iii) their leadership qualities; and
- iv) the foundations for future accomplishments that they laid down and their ongoing influence upon later generations.

I shall deal with them chronologically.

Robert Falcon Scott led two expeditions, in 1901-04 and in 1910-13. Both were in the Ross Sea sector. His exploration forays were only 300 miles in extent, according to the historian J. Gordon Hayes (1928). However, he and the men of his two expeditions discovered about 1300 miles of coast, mainly in the N.Z. sector.

Note how little was explored. It is comparatively easy to discover features. You can, for example, sail along a coast, 40 miles off-shore, and make notes of mountains, bays and capes that you observe.

Exploration is more difficult. You must land and survey the features, accurately determining their exact positions (generally by astronomical observations).

Scott's scientific results were extensive, with many volumes of scientific reports covering a variety of disciplines. They were uniquely valuable because in many subjects they provided the first information ever gained in the Antarctic and opened up various avenues for research by later expeditions. The high standards achieved were due to the quality of the scientists that were included in his teams.

However, despite these accomplishments, Scott

has been widely criticized. Errors in planning, combined with rare bad luck, led to the deaths of the five men in his South Pole party. He was autocratic and reluctant to accept advice or to consult others. He was an amateur who lacked experience in field work and made no attempts to prepare himself for Antarctic conditions either by previous activity in the frigid regions of Europe or research of available literature.

During some eighty-odd years of searching for the NW passage and a route to the North Pole, the Royal Navy had despatched more than 30 expeditions. The experience gained from these covered man-hauling, dog-sledging, food, scurvy, and survival procedures. Captain Scott, R.N., did not bother to read the extensive material available in the Royal Navy archives and repeated many of the mistakes that these early Arctic explorers had made.

Yet the tragic story of Scott's fate, with its engrossing examples of endurance and fortitude, did more to stimulate adventure in young people and later Antarctic endeavour than probably any other narrative in the British language. And the emotion that his fate evoked led to an appeal for funds that resulted in support for later explorers and, more important, to the establishment of the Scott Polar Research Institute in Cambridge, a world-renowned institute.

So, all in all, I rate Scott very highly overall, but not for his personal exploration, which was not very much.

The Shackleton saga is quite a different one. Contrary to general opinion, Shackleton's life was a succession of failures if regarded from the point of view of his objectives. He accompanied Scott on his first attempt to reach the South Pole in 1903, and, stricken with scurvy, spent part of their retreat being carried on a sledge. He was shipped back to England and missed the second year of the expedition.

On Shackleton's 1907-09 expedition he was forced to abandon his own attempt to reach the Pole when, according to his records, he was within 96 miles of it.

His third expedition, 1914-17, failed when his ship *Endurance* was crushed by pack ice before he had reached the Continent. And on his final expedition, 1921-22, he died at South Georgia before proceeding into Antarctic waters.

In addition to all this, Shackleton embarked upon a number of get-rich-quick gold mining financial ventures, each of which failed.

He, himself, explored very little, his main achievement being pioneering the Pole route via the Beardmore Glacier. He explored a stretch of terrain



about 200 miles in extent during his 1907-09 expedition.

His men, however, achieved much more. David, Mawson and McKay made a sledge journey of 1200-odd miles to reach the South Magnetic Pole, during which Mawson surveyed in detail about 200 miles of the west coast of the Ross Sea. Shackleton, in his ship, explored the edge of the Ross Ice Shelf. Hayes awards Shackleton 300 miles of exploration and his expedition 1035 miles of discovery.

Scientifically, Shackleton's 1907-09 expedition was productive for, like Scott, he had a fine team of scientists. The output for this one expedition was somewhat less than half of that of Scott's two expeditions.

Shackleton was a democratic leader, in an era when leadership was almost universally autocratic, and this, with his extrovert and magnetic personality, gained him the loyalty and affection of his men. As a leader in the field, he was outstanding.

The Shackleton story, like Scott's, had a profound effect upon generations of young and old, his survival and his leadership, following the destruction of his ship *Endurance*, being one of the greatest narratives of British adventure. It is largely upon this that his fame rests, and for this he scores well under criterion four.

As an interesting footnote to all this, Sir Douglas Mawson once told me that he believed Shackleton faked his travel records and that he could not have reached a point 96 miles from the Pole. Mawson based his judgement upon his own experience of man-hauling a sledge for 1200 miles. The "Grey Eminence" of 19<sup>th</sup> Century British polar exploration, Sir Clement Markham, expressed the same view, challenging the daily distances traversed, particularly in ascending the Beardmore Glacier.

Several years ago in England, a descendant of a man who accompanied Shackleton on this sledge journey, released the diary of this man, which specifically states that the Boss, as he was called, cribbed a few miles each day when he entered up his travel log!

All in all, I rate Shackleton as a very great adventurer, but not much of an explorer, and I do not give him nearly as many marks as Scott.

Now consider Amundsen. Whereas Scott and Shackleton were amateurs, Amundsen was highly professional. He had spent twenty or more years preparing himself for his conquest of the South Pole. He had gained Antarctic experience when, as second mate, he wintered on De Gerlache's ship, the *Belgica*, which was beset for the winter in Antarctica in 1898.

He studied for, and gained, his Master's ticket and was experienced in skiing and dog sledging. Further, he had carefully studied the exploits of that great Polar traveller, Fridtjof Nansen.

Amundsen's route to the Pole, via the Axel Heiberg Glacier, was steeper, more crevassed and much more difficult than that of Scott up the Beardmore, as Wally Herbert, a British expeditioner, demonstrated, shortly after the IGY, when he drove a dog team down it. Yet Amundsen reached his goal and returned with a minimum of fuss. The main criticism of his expedition is that he did no exploration, i.e. surveying, as he went along, so little in fact, that it is even difficult to plot accurately the route he took. The real exploration of the Axel Heiberg Glacier and its environs was not carried out until the IGY, more than 50 years later.

Nor did Amundsen's party carry out any scientific work. His journey was essentially a rapid dash to reach a defined goal, nothing more. Hayes says of it: "It practically was wasted work, because no proper cartographical record was made, even of the bare route; whereas there was ample (dog) power, food and time to have explored the Queen Maud Range and Carmen Land. Amundsen should have taken at least one competent surveyor on his Polar journey; for it is really no use breaking the first trail unless something tangible is done to open up the new country".

Hayes quotes the extent of new land discovered (but not explored) as 1300 miles, although Amundsen logged 1612 miles of actual sledging.

It is ironical that, in the Arctic, Amundsen's record as an adventurer, a discoverer and an explorer is at the highest level; it was he who discovered – and charted – the North-West Passage!

Even he had his critics when it came to leadership. Loyal to his friends, he could be quite vindictive if opposed. When returning to his base from an abortive early attempt on the South Pole route, which he began too early in the Spring, he deserted his party and raced on ahead. As a result, two sledge groups nearly failed to reach base. One man, Johannsen, bitterly rebuked him after returning. Amundsen regarded this as mutiny, removed him from the final assault party, and ostracized him for the rest of the expedition. Shamed and humiliated, the man later, in Norway, committed suicide.

Rated accordingly to my four criteria, Amundsen does not score a high total in Antarctic endeavour, although like Shackleton, he gets high marks as an adventurer.

Moving on now to Mawson, we find a large mile-



age for personal exploration if we include his long magnetic pole journey while serving with Shackleton. In his AAE and BANZARE expeditions, the total for personal exploration is only about 300 miles, for on his second expedition, BANZARE, he achieved practically no personal exploration. He made only three landings in two summer voyages and made no cartographic observations on any of them.

In miles of coast discovered, he rates far more highly, with around 1800 miles to his credit, for his BANZARE voyages sighted stretches of previously unknown coast while passing by, out to sea.

Mawson's scientific results were extensive, with observations in various disciplines from Macquarie Island, Commonwealth Bay, the Western Base on the Shackleton Ice Shelf, and marine biology and oceanography during the BANZARE voyages.

As a leader, Mawson rates more highly than Scott and Amundsen. First, his style was more democratic and, second, he was himself a scientist and thus was able to relate better to the scientists under his command. Also, he personally ensured a high priority for all the scientific work and worked hard later to see that the results were published.

With reference to criterion number four, it can be said that Mawson's influence was great. His work has been an inspiration to numerous young adventurers in Australia while, in an ongoing role, his was the major influence in persuading the Commonwealth Government in 1947 to establish the ANARE. In the early days of my expeditions, while serving on the ANARE Executive Planning Committee, he was a powerful supporter of my plans and programs.

Hayes has this to say of him :

"Sir Douglas Mawson's 1911-13 expedition may be held up as a model for others to copy. Its excellence lay in its design, its scope and its executive success; and it owes its exalted position among other Antarctic expeditions mainly to the fact that it was originated and conducted by scientists of administrative ability who are the fit and proper persons for such undertakings. Reason appears to have ruled throughout."

I have also applied my criteria to other Antarctic leaders. I should have liked to discuss Richard Byrd (USA), Finn Ronne (USA), John Rymill (Australia), Vivian Fuchs (UK), Edmund Hillary (NZ), Mikhail Somov (Russia) – for no analysis of Antarctic endeavour would be adequate without examining their work, but time does not permit.

You should understand that the four expeditions I have treated provided merely an introduction to the

opening up of the last unknown continent. Extensive exploration of Antarctica did not occur until the years 1950-70, particularly during the International Geophysical Year (1957-58) and the years that followed. Australians figured prominently in this work, discovering vast tracts of unknown country and exploring in detail more than a million square kilometres of terrain. One ANARE man, surveyor Syd Kirkby, has explored more than the explorations of Scott, Shackleton, Amundsen and Mawson combined! His is a remarkable record.

He was the wintering surveyor at Mawson in 1956 and 1960 and returned in 1980 as Station Leader and surveyor. He accompanied me on summer voyages of exploration in 1955-56, 1956-57, 1960-61 and 1962-63. He was the hands-on explorer, the man who actually surveyed and mapped the country.

His surface exploration – by dog sledges, weasels and tractor trains – covered 1200 km of "new" territory. His actual travel distances are much greater. The lengths of coastline he explored with me, i.e. coasts where no-one had been before, totalled 3000 km.

By air, during exploration flights and flights to achieve first landings, he covered approximately 3000 km. His astrofixes range from 44 degrees E to 167 degrees E longitude and inland to the most southerly nunataks of the Prince Charles Mountains.

No other person, living or dead, has done so much!

The IGY and the years that followed produced accomplishments by a number of remarkable men whose exploits have received no wide acclaim. I remember particularly Americans Crary and Bentley and Russians Somov and Kapitsa. I have been urging for years that more comprehensive histories of Antarctic exploration should be carried out, dealing with the major explorations of the years 1950-70 and omitting the work of the heroic era, which has been rehashed time after time and is still attracting the attention of book authors.

My own judgement, ranking all the men I have mentioned, according to the criteria I stated at the beginning, is that Scott, Mawson, Byrd and Fuchs are well ahead of the rest. However, what is needed is a modern J. Gordon Hayes to survey the whole panorama of Antarctic history in detail. Only then will Antarctic exploration be presented to public view in its true perspective.









PROCEEDINGS  
OF THE  
ROYAL SOCIETY OF VICTORIA  
INCLUDING  
TRANSACTIONS OF MEETINGS

Volume 115

NUMBERS 1/2

ROYAL SOCIETY'S HALL  
9 VICTORIA STREET, MELBOURNE, VICTORIA 3000



## CONTENTS OF VOLUME 115, NUMBER 1/2

### PROCEEDINGS

- B. MACKNESS 1  
A review of nematophagy by terrestrial tardigrades with new observations on some Australian species.
- BASDEN, ALISON 7  
Early Devonian (Emsian) vertebrate microremains from the Buchan Group, Victoria, Australia
- FRAMENAU, VOLKER W. 27  
Two Alpine Wolf Spiders of Australia: *Artoria alta* sp. nov., and the male of *Lycosa musgravei* McKay, 1974 (Araneae, Lycosidae)
- GERDTZ, W. R. & ARCHBOLD, N. W. 35  
*Glaucodon ballaratensis* (Marsupialia, Dasyuridae), a late Pliocene 'devil' from Batesford, Victoria
- GERDTZ, W. R. & ARCHBOLD, N. W. 45  
An early occurrence of *Sarcophilus lanianus harrisii* (Marsupialia, Dasyuridae) from Nelson Bay, Victoria
- YEN, ALAN L. & WAINER, JOHN 55  
Scolopendrid centipedes caught by pitfall trapping in the Uluru-Kata Tjuta National Park, Northern Territory, including a record of an apparent 'outbreak'
- BIRCH, W. D. & SAMUELS, L. E. 67  
The Ballarat meteorite, a fossil IAB iron from Victoria, Australia

### TRANSACTIONS

- LAW, PHILLIP i  
Lecture to the Danish Club, 18<sup>th</sup> September, 2003



# ROYAL SOCIETY OF VICTORIA

2003

<i>Patron:</i>	HIS EXCELLENCY, THE HONORABLE JOHN LANDY, AC, MBE GOVERNOR OF VICTORIA
<i>President:</i>	N. W. ARCHBOLD, BA, MSc, PhD
<i>Vice-President:</i>	B. G. LIVETT, BSc(HONS), PhD
<i>Immediate Past President:</i>	G. D. SANSON, BSc(HONS), PhD
<i>Honorary Secretary:</i>	J. W. WARREN, MA, PhD
<i>Honorary Treasurer:</i>	B. J. WALBY, MA, MSc, CChem, MRSC, FIMA
<i>Honorary Librarian:</i>	D. A. McCANN, PhD
<i>Honorary Editor:</i>	M. GROVER, BEd, BSc(HONS)
<i>Honorary Research Secretary:</i>	W. D. BIRCH, BSc(HONS), PhD

The above Office-bearers are *ex officio* members of the Council of the Society.

Other members of the Council are:

P. G. BAINES, BA(HONS), BSc, PhD  
J. M. BOWLER, AM, MSc, PhD, FAHA, FRSV  
R. J. FOSTER, BE  
M. R. HARRIS, BSc(HONS)  
J. F. LOVERING, MSc, PhD, FAA, FTS  
W. J. W. MCAULEY, MSc, MAIMM, FAIE, FRGS, FRNSW, MAIG, MGSA  
P. C. MANINS, BSc, BE(HONS), PhD, CCM, QEP, PSM, FTSE  
N. F. MILLIS, AC, MBE, MAGrSc, PhD, FTSE, FRSV  
L. SELWOOD, MSc, PhD  
D.M. STOKES, BSc(HONS), PhD  
P. G. THORNE, BSc, PhD, DIPPubPOL  
M. WILLIAMS, BSc, MS

*Executive Officer:*

C. B. VAN MEGEN

<i>Honorary Financial Advisor:</i>	FOSTER STREADER BLACK PTY LTD
<i>Honorary Solicitors:</i>	PHILLIPS FOX
<i>Honorary Auditor:</i>	M. KOVASSY, MAcc, CPA, AAIM
<i>Trustees:</i>	D. M. CHURCHILL, BSc, MA, AAIM, PhD PROFESSOR J. W. WARREN, MA, PhD HON. SIR EDWARD WOODWARD, KT, OBE, LL.M., QC

## INSTRUCTIONS FOR AUTHORS

Papers considered for publication may be: Reviews, Reports of experimental or descriptive research, or Short Communications. Length of papers may vary but Short Communications should not exceed 1500 words. When an author submits a manuscript for consideration, it is assumed that it has never been published or is under consideration for publication elsewhere.

### SUBMISSION

The original and two copies of the typescript and all tables and figures should be submitted to the Executive Officer, Royal Society of Victoria, 9 Victoria Street, Melbourne, Victoria, 3000. Use International A4 bond paper, printed on one side only. In a letter of transmittal, give the names and addresses (postal and email) of the author or co-authors, also give the names, addresses and email addresses of three persons not directly associated with the work and outside the author's institution, who could act as referees. For all papers, presentation of the final manuscript on word-processed floppy disk is essential as well as a hard copy. Authors or their institutions may be requested to contribute towards the cost of publication of the paper and in the case of very long papers, such contribution is mandatory.

It will assist reviewers if related papers recently published or submitted elsewhere accompany the submitted manuscript.

### FORMAT

All manuscripts should be written in clear and concise English. Use double spacing throughout; leave 30 mm margins around the text; number all pages. All measurements are to be expressed in SI units (eg  $\mu\text{m}$ , mm, m, km but not cm) and standard symbols and abbreviations used. Note, Ma refers to a date (eg 25Ma) not a time interval, which is written as 5 million years. When a number does not refer to a measurement it should be spelt out except when it is greater than nine. Wherever possible fractions should be written in the form x/y.

Geological papers must follow the Australian Code of Stratigraphic Nomenclature and should cite geological maps in the text with italics for the 1:63 360 series and full caps for the 1:250 000 series. New stratigraphic names should be registered with the Geoscience Australia (GA) Australian Stratigraphic Names Database. The GA website "How to define a lithostratigraphic unit" ([http://www.ga.gov.au/minerals/strat\\_names/](http://www.ga.gov.au/minerals/strat_names/)) is a useful guide.

The Proceedings uses Australian spelling according to the latest edition of the *Macquarie Dictionary* and where there is optional spelling for words we use the form used by Victorians traditionally. For example we use the following spelling; color, honorary, program.

Authors should follow the layout of headings, tables and illustrations as presented in a recent issue of the *Proceedings*.

Papers should be organised as follows:

1. *A brief title*, capitalised and, if possible, beginning with a key word.
2. *The name and address of the author(s)*, with numerical superscripts to distinguish addresses of multiple authors.
3. *A full reference to the paper*, leaving space for the printer's additions.
4. *A short abstract* of not more than 200 words describing the results (rather than the contents) of the paper.
5. *Up to 5 key words*.
6. *The main text*. Capitalise the first word of the introductory paragraph; do not use the heading 'Introduction'. Within the text up to three grades of headings may be used, typed as follows:

#### GRADE ONE HEADING

*Grade two heading*

*Grade three heading*. Followed by running text on the same line.

Refer to illustrations of the paper in the text as 'Fig. 1A, B', 'Figs 1, 2' or 'Figs 1-4', and indicate in the margins where the illustrations should be placed. Refer to references in the text as Archbold (1998), (Archbold 1998) or (Archbold 1998: 2, fig. 1); multiple citations of references should be arranged chronologically. All references cited in the text must be listed at the end of the paper. Footnotes in the main text are not allowed. Where there are multiple authors for a reference, "et al." is used, and is not italicized.

In taxonomic works, synonymies should be of the same format as the following examples, with a dash preceding authors' names except in the case of reference to the original description.

*Endendrium generalis* Lendenberg 1885: 351, pl. 6. — Lendenberg 1887: 16.

*Endendrium generale*. — Hartlaub 1905: 515.—Watson 1985: 196-200, figs 40-52.

non *Endendrium generale*.—Watson 1982: 89, pl. 10, fig. 3.

*Endendrium lendenfeldi* Briggs 1922: 150.—Rosler 1978: 104, 120, pl. 20, figs 1-3.

Note that plate and figure numbers, etc. originally given in Roman numerals should be transliterated into Arabic figures; this is also the case in the main text and in the references.

7. *Acknowledgements*. The source of financial grants and other funding as well as the contribution of colleagues or institutions should be acknowledged. These should follow the main text and be as brief as possible.

8. *References*. These should conform in arrangement to the examples below. Journal titles must be cited in full as they appear on the title page. No abbreviations will be accepted unless they are standard abbreviations such as "Co." for company, or "ed." for editor, "eds" for editors..

CURTIS, N. P., 2001. Germination of *Xanthorrhoea anstralis* using treatments that mimic post-fire and unburnt conditions. *Proceedings of the Royal Society of Victoria* 113(2): 237-245.

BERGSON, H., 1928. *Creative Evolution*. MacMillan and Co., Limited, London, xv+425 pp.

ROSEN, B.R. & TURNŠEK, D., 1989. Extinction patterns and biogeography of scleraetinian corals across the Cretaceous/Tertiary boundary. In *Proceedings of the Fifth International Symposium on Fossil Cnidaria including Archaeocyatha and Spongionomorphs*, P.A. Jell & J.W. Pickett, eds, Association of Australasian Palaeontologists, Brisbane, Queensland, 355-370.

Personal communications and unpublished data are not to be listed in the reference list but should be mentioned in full in the text (eg P. Brown pers. comm. 2002).

9. *Tables and Figures*. Each table with its title should be typed on a separate sheet. A separate sheet should also be used to list captions to figures in numerical order.

### ILLUSTRATION

These must be designed for one or two column width (67 mm or 140 mm); maximum length is 193 mm. Composite Figures should be labelled A, B, C, etc. (not a, b, c). Oversized illustrations, tables or maps may be accepted for publication as foldouts only if the author meets all costs involved in their production. Maximum size for foldouts is 193x285 mm.

Final submission of black and white figures electronically is encouraged as long as it is in a readily acceptable format (JPEG, TIFF, emf, wmf, pcx, pct, wpg, cdr). These can be

supplied on 3.5" floppy disks, zip disks or CD (especially for figures with any grey shades) and accompanied by a high quality hard copy. Digital files must be supplied at a resolution of at least 600 d.p.i. at full size.

### PROOFS

Authors will normally receive an 'Adobe Acrobat PDF' version of the mock page proofs; this is for the purpose of correcting printer's errors and not for altering the wording or substance of the paper. This should be returned to the editor within 72 hours. If absent, authors should arrange for a colleague to check and return proofs to the editor on their behalf. Authors will be charged for excessive alterations at this stage.

### REPRINTS

The corresponding author will be emailed an 'Adobe Acrobat PDF' of the final printed manuscript at the time of printing. Hardcopy reprints may be ordered by returning the order form that accompanies the mock page proofs.

Upon publication, copyright of the paper (including illustrations) is held by the Royal Society of Victoria. Any subsequent publication of any material shall be by written permission of the Society. Authors in Government institutions where copyright is retained by the Crown are exempt from transfer of copyright to the Society.





51321—





# PROCEEDINGS OF THE ROYAL SOCIETY OF VICTORIA

Volume 115, Numbers 1/2

## CONTENTS

### PROCEEDINGS

- B. MACKNESS 1  
A review of nematophagy by terrestrial tardigrades with new observations on some Australian species.
- BASDEN, ALISON 7  
Early Devonian (Emsian) vertebrate microremains from the Buchan Group, Victoria, Australia
- FRAMENAU, VOLKER W. 27  
Two Alpine Wolf Spiders of Australia: *Artoria alia* sp. nov., and the male of *Lycosa musgravei* McKay, 1974 (Araneae, Lycosidae)
- GERDTZ, W. R. & ARCHBOLD, N. W. 35  
*Glaucodon ballaratensis* (Marsupialia, Dasyuridae), a late Pliocene 'devil' from Batesford, Victoria
- GERDTZ, W. R. & ARCHBOLD, N. W. 45  
An early occurrence of *Sarcophilus laniarius harrisii* (Marsupialia, Dasyuridae) from Nelson Bay, Victoria
- YEN, ALAN L. & WAINER, JOHN 55  
Scolopendrid centipedes caught by pitfall trapping in the Uluru-Kata Tjuta National Park, Northern Territory, including a record of an apparent 'outbreak'
- BIRCH, W. D. & SAMUELS, L. E. 67  
The Ballarat meteorite, a fossil IAB iron from Victoria, Australia

### TRANSACTIONS

- LAW, PHILLIP i  
Lecture to the Danish Club, 18<sup>th</sup> September, 2003